

On the Karaś type theorems for the multidegrees of polynomial automorphisms

Shigeru Kuroda

Abstract

To solve Nagata's conjecture, Shestakov-Umirbaev constructed a theory for deciding wildness of polynomial automorphisms in three variables. Recently, Karaś and others study multidegrees of polynomial automorphisms as an application of this theory. They give various necessary conditions for triples of positive integers to be multidegrees of tame automorphisms in three variables. In this paper, we prove a strong theorem unifying these results using the generalized Shestakov-Umirbaev theory.

1 Introduction

Let k be a field of characteristic zero, and $k[\mathbf{x}] = k[x_1, \dots, x_n]$ the polynomial ring in n variables over k . In this paper, we discuss the group $\text{Aut}_k k[\mathbf{x}]$ of automorphisms of the k -algebra $k[\mathbf{x}]$. To denote an element of $\text{Aut}_k k[\mathbf{x}]$, we often use the notation $F = (f_1, \dots, f_n)$, where each f_i represents the image of x_i under the automorphism. We say that F is *elementary* if there exist $1 \leq l \leq n$, $a \in k^\times$ and $p \in k[\mathbf{x} \setminus \{x_l\}]$ such that $f_l = ax_l + p$ and $f_i = x_i$ for each $i \neq l$, where we regard $\mathbf{x}$ as the set $\{x_1, \dots, x_n\}$ of variables. The subgroup $T_n(k)$ of $\text{Aut}_k k[\mathbf{x}]$ generated by all the elementary automorphisms is called the *tame subgroup*. We say that F is *tame* if F belongs to $T_n(k)$, and *wild* otherwise.

The following problem is one of the fundamental problems in Polynomial Ring Theory.

Problem 1.1. *Does it hold that $\text{Aut}_k k[\mathbf{x}] = T_n(k)$?*

2010 *Mathematical Subject Classification.* Primary 14R10; Secondary 13F20, 16Z05.

Partly supported by the Grant-in-Aid for Young Scientists (B) 24740022, Japan Society for the Promotion of Science.

Since k is a field, every automorphism of $k[\mathbf{x}]$ is elementary if $n = 1$. Due to Jung [3], $\text{Aut}_k k[\mathbf{x}] = T_2(k)$ holds for $n = 2$. In the case of $n = 3$, Nagata [17] conjectured that $\text{Aut}_k k[\mathbf{x}] \neq T_3(k)$, and constructed a candidate of wild automorphism. Recently, Shestakov-Umirbaev [18], [19] solved the conjecture in the affirmative. They not only proved that Nagata's automorphism is wild, but also gave a criterion for deciding wildness of elements of $\text{Aut}_k k[\mathbf{x}]$ when $n = 3$. Problem 1.1 remains open for $n \geq 4$.

Now, for each $F \in \text{Aut}_k k[\mathbf{x}]$, we define the *multidegree* of F by

$$\text{mdeg } F = (\deg f_1, \dots, \deg f_n).$$

Here, $\deg f$ denotes the total degree of f for each $f \in k[\mathbf{x}]$. We set

$$\text{mdeg } S = \{\text{mdeg } F \mid F \in S\}$$

for each subset S of $\text{Aut}_k k[\mathbf{x}]$.

The following proposition is due to Karaś [5, Proposition 2.2].


Proposition 1.2 (Karaś). *Let $d_1, \dots, d_n \in \mathbf{N}$ be such that $d_1 \leq \dots \leq d_n$. If there exists $2 \leq i \leq n$ such that d_i belongs to*

$$\mathbf{N}_0 d_1 + \dots + \mathbf{N}_0 d_{i-1} := \{u_1 d_1 + \dots + u_{i-1} d_{i-1} \mid u_1, \dots, u_{i-1} \in \mathbf{N}_0\},$$

then $(d_1, \dots, d_n)$ belongs to $\text{mdeg } T_n(k)$.

Here, $\mathbf{N}$ and $\mathbf{N}_0$ denote the sets of positive integers and nonnegative integers, respectively. In the same paper, Karaś proved that $(3, 4, 5)$, $(3, 5, 7)$, $(4, 5, 7)$ or $(4, 5, 11)$ do not belong to $\text{mdeg } T_3(k)$ using the theory of Shestakov-Umirbaev [18], [19]. This means that any automorphism with multidegree $(3, 4, 5)$, $(3, 5, 7)$, $(4, 5, 7)$ or $(4, 5, 11)$ is wild.

Following this paper, Karaś and others recently study detailed conditions for elements of $\mathbf{N}^3$ to belong to $\text{mdeg } T_3(k)$. They derive from the Shestakov-Umirbaev theory various conditions under which $(d_1, d_2, d_3) \in \mathbf{N}^3$ belongs to $\text{mdeg } T_3(k)$ if and only if d_2 is divisible by d_1 , or d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$. Because of Proposition 1.2, this is the same as giving conditions under which $(d_1, d_2, d_3) \in \mathbf{N}^3$ does not belong to $\text{mdeg } T_3(k)$ if d_2 is not divisible by d_1 , and d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$. By a *Karaś type theorem*, we mean a theorem which describes such a condition. In the following diagram, the oval represents the set of elements of $\mathbf{N}^3$ satisfying one of such conditions.



Since many partial results were published separately, we consider it necessary to prove a strong theorem unifying them. The purpose of this paper is to interpret the generalized Shestakov-Umirbaev theory [11], [12] in terms of multidegrees. This naturally leads to a very general version of Karaš type theorem (Theorem 2.5) which is also valid for the case of “weighted” degree. In the case of total degree, our result implies the following theorem.

For $d_1, d_2, d_3 \in \mathbf{N}$, consider the conditions (a), (b) and (c) as follows:

- (a) One of the following holds:
 - (a1) $3d_2 \neq 2d_3$ and $sd_1 \neq 2d_3$ for any odd number $s \geq 3$.
 - (a2) $d_1 + d_2 \leq d_3 + 2$.
- (b) One of the following holds:
 - (b1) $\gcd(d_1, d_2, d_3) = \gcd(d_1, d_2) \leq 3$.
 - (b2) $d_1 + d_2 + d_3 \leq \text{lcm}(d_1, d_2) + 2$.
- (c) d_2 is not divisible by d_1 , and d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.

Then, we have the following theorem.

Theorem 1.3. *Let $d_1 \leq d_2 \leq d_3$ be positive integers satisfying (a), (b) and (c). Then, (d_1, d_2, d_3) does not belong to $\text{mdeg } T_3(k)$.*

We note that (a) holds if and only if (a1) or one of the following holds:

- (a2') $3d_2 = 2d_3$ and $2d_1 \leq d_2 + 4$.
- (a2'') $sd_1 = 2d_3$ and $2d_2 \leq (s - 2)d_1 + 4$ for some odd number $s \geq 3$.

Actually, (a2) is equivalent to $2d_1 \leq d_2 + 4$ if $3d_2 = 2d_3$, and to $2d_2 \leq (s - 2)d_1 + 4$ if $sd_1 = 2d_3$ for some odd number $s \geq 3$. The condition (b1) is

equivalent to $\gcd(d_1, d_2) \leq 3$ and $\gcd(d_1, d_2) \mid d_3$, which is fulfilled whenever $\gcd(d_1, d_2) = 1$. By (c), Theorem 1.3 is considered as a Karaš type theorem. As will be discussed in Section 6, this theorem implies all the Karaš type theorems which the author knows, except for Karaš [8, Theorem 1]. This result of Karaš can be derived from Theorem 2.5 by assuming the lower bound for the degrees of “Poisson brackets” given in [8, Theorem 14].

It is easy to see that, when $n \geq 3$, there exists $(d_1, \dots, d_n) \in \text{mdeg } T_n(k)$ with $d_1 \leq \dots \leq d_n$ for which d_i does not belong to $\mathbf{N}_0 d_1 + \dots + \mathbf{N}_0 d_{i-1}$ for any $2 \leq i \leq n$. For instance, take any prime numbers $p_1 < \dots < p_{n-1}$, and define $d_i = p_i^i p_{i+1} \dots p_{n-1}$ for each $i \neq n$, and $d_n = (d_{n-1} - 1)d_{n-2} + 1$. Then, we have $d_1 \leq \dots \leq d_n$, and d_i does not belong to $\mathbf{N}_0 d_1 + \dots + \mathbf{N}_0 d_{i-1}$ for any $2 \leq i \leq n$, since $p_{i-1} \nmid d_i$. Moreover, $(d_1, \dots, d_n)$ is the multidegree of $F \in T_n(k)$ defined by $f_i = x_i + x_n^{d_i}$ for each $i \neq n$, and $f_n = x_n + f_{n-2}^{d_{n-1}} - f_{n-1}^{d_{n-2}}$. Thus, it is also an interesting problem to find a good *sufficient* condition for elements of $\mathbf{N}^n$ to belong to $\text{mdeg } T_n(k)$. We mention that the situation is more complicated in the case of weighted degree. In this case, a statement similar to Proposition 1.2 does not hold in general. It seems difficult to expect a simple criterion as in the case of total degree.

The structure of this paper is as follows. In Section 2, we state the main theorem after recalling basic definitions. Sections 3, 4 and 5 are devoted to the proof of the main theorem. In Section 3, we prove a key theorem using the generalized Shestakov-Umirbaev theory. In Section 4, we generalize Sun-Chen [20, Lemma 3.1] to the case of weighted degree. The proof of the main theorem is completed in Section 5 using the results in Sections 3 and 4. In Section 6, we prove a variety of Karaš type theorems as applications of our theorems.

2 Main results

First, we recall basics of the weighted degrees of polynomial automorphisms and differential forms. Let Γ be a *totally ordered additive group*, i.e., an additive group equipped with a total ordering such that $\alpha \leq \beta$ implies $\alpha + \gamma \leq \beta + \gamma$ for each $\alpha, \beta, \gamma \in \Gamma$. We denote by Γ_+ the set of positive elements of Γ . Let $\mathbf{w} = (w_1, \dots, w_n)$ be an n -tuple of elements of Γ_+ . We define the **w-weighted Γ -grading**

$$k[\mathbf{x}] = \bigoplus_{\gamma \in \Gamma} k[\mathbf{x}]_\gamma$$

by setting $k[\mathbf{x}]_\gamma$ to be the k -vector subspace of $k[\mathbf{x}]$ generated by $x_1^{a_1} \dots x_n^{a_n}$ for $a_1, \dots, a_n \in \mathbf{N}_0$ with $\sum_{i=1}^n a_i w_i = \gamma$ for each $\gamma \in \Gamma$. Write $f \in k[\mathbf{x}]$ as $f = \sum_{\gamma \in \Gamma} f_\gamma$, where $f_\gamma \in k[\mathbf{x}]_\gamma$ for each $\gamma \in \Gamma$. Then, we define the **w-degree**

of f by

$$\deg_{\mathbf{w}} f = \max\{\gamma \in \Gamma \mid f_{\gamma} \neq 0\}.$$

Here, the maximum of the empty set is defined to be $-\infty$. We define $f^{\mathbf{w}} = f_{\delta}$ if $\delta := \deg_{\mathbf{w}} f > -\infty$, and $f^{\mathbf{w}} = 0$ otherwise. Then, we have

$$\deg_{\mathbf{w}} fg = \deg_{\mathbf{w}} f + \deg_{\mathbf{w}} g \quad \text{and} \quad (fg)^{\mathbf{w}} = f^{\mathbf{w}} g^{\mathbf{w}} \quad (2.1)$$

for each $f, g \in k[\mathbf{x}]$. If $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, \dots, 1)$, then the $\mathbf{w}$ -degree of f is the same as the total degree of f .

We remark that, if $\deg_{\mathbf{w}} f < w_i$ for $f \in k[\mathbf{x}]$ and $1 \leq i \leq n$, then f belongs to $k[\mathbf{x} \setminus \{x_i\}]$. As a consequence of this fact, we obtain the following lemma.

Lemma 2.1. *Let $\mathbf{w} \in (\Gamma_+)^n$ and $F \in \text{Aut}_k k[\mathbf{x}]$ be such that*

$$w_1 \leq \dots \leq w_n \quad \text{and} \quad \deg_{\mathbf{w}} f_1 \leq \dots \leq \deg_{\mathbf{w}} f_n.$$

Then, we have $\deg_{\mathbf{w}} f_i \geq w_i$ for $i = 1, \dots, n$.

Proof. Suppose to the contrary that $\deg_{\mathbf{w}} f_i < w_i$ for some $1 \leq i \leq n$. Then, we have $\deg_{\mathbf{w}} f_j < w_l$ for each $1 \leq j \leq i$ and $i \leq l \leq n$. This implies that $f_1, \dots, f_i$ belong to $k[x_1, \dots, x_{i-1}]$. Hence, $f_1, \dots, f_i$ are algebraically dependent over k , a contradiction. $\square$

For each $F \in \text{Aut}_k k[\mathbf{x}]$, we define the $\mathbf{w}$ -degree and $\mathbf{w}$ -multidegree by

$$\deg_{\mathbf{w}} F = \sum_{i=1}^n \deg_{\mathbf{w}} f_i \quad \text{and} \quad \text{mdeg}_{\mathbf{w}} F = (\deg_{\mathbf{w}} f_1, \dots, \deg_{\mathbf{w}} f_n),$$

respectively. For each subset S of $\text{Aut}_k k[\mathbf{x}]$, we define

$$\text{mdeg}_{\mathbf{w}} S = \{\text{mdeg}_{\mathbf{w}} F \mid F \in S\}.$$

Set $|\mathbf{w}| = w_1 + \dots + w_n$. Then, we know by Lemma 2.1 that $\deg_{\mathbf{w}} F \geq |\mathbf{w}|$, and $\deg_{\mathbf{w}} F = |\mathbf{w}|$ if and only if $\text{mdeg}_{\mathbf{w}} F = (w_{\sigma(1)}, \dots, w_{\sigma(n)})$ for some $\sigma \in \mathfrak{S}_n$. Here, $\mathfrak{S}_n$ denotes the symmetric group of $\{1, \dots, n\}$. Thus, we may assume that $\deg_{\mathbf{w}} F > |\mathbf{w}|$ in the study of $\text{mdeg}_{\mathbf{w}} F$. We mention that F is tame if $\deg_{\mathbf{w}} F = |\mathbf{w}|$ for some $\mathbf{w} \in (\Gamma_+)^n$ by [12, Lemma 6.1 (i)].

The notion of the $\mathbf{w}$ -degrees of differential forms is important in the study of polynomial automorphisms. Let $\Omega_{k[\mathbf{x}]/k}$ be the module of differentials of $k[\mathbf{x}]$ over k , and $\bigwedge^r \Omega_{k[\mathbf{x}]/k}$ the r -th exterior power of the $k[\mathbf{x}]$ -module $\Omega_{k[\mathbf{x}]/k}$ for $r \in \mathbf{N}$. Then, each $\omega \in \bigwedge^r \Omega_{k[\mathbf{x}]/k}$ is uniquely expressed as

$$\omega = \sum_{1 \leq i_1 < \dots < i_r \leq n} f_{i_1, \dots, i_r} dx_{i_1} \wedge \dots \wedge dx_{i_r},$$

where $f_{i_1, \dots, i_r} \in k[\mathbf{x}]$ for each $i_1, \dots, i_r$. Here, df denotes the differential of f for each $f \in k[\mathbf{x}]$. We define the $\mathbf{w}$ -degree of ω by

$$\deg_{\mathbf{w}} \omega = \max\{\deg_{\mathbf{w}} f_{i_1, \dots, i_r} x_{i_1} \cdots x_{i_r} \mid 1 \leq i_1 < \cdots < i_r \leq n\}.$$

When $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, \dots, 1)$, we denote $\deg_{\mathbf{w}} \omega$ simply by $\deg \omega$. By definition, we have

$$\deg_{\mathbf{w}} df \wedge dg = \max\left\{\deg_{\mathbf{w}} \left(\frac{\partial f}{\partial x_i} \frac{\partial g}{\partial x_j} - \frac{\partial f}{\partial x_j} \frac{\partial g}{\partial x_i}\right) x_i x_j \mid 1 \leq i < j \leq n\right\}$$

for $f, g \in k[\mathbf{x}]$. Hence, $\deg df \wedge dg$ is the same as the degree of the *Poisson bracket* $[f, g]$ defined by Shestakov-Umirbaev [18].

For each $F \in \text{Aut}_k k[\mathbf{x}]$, we have $df_1 \wedge \cdots \wedge df_n = c dx_1 \wedge \cdots \wedge dx_n$, where $c \in k^\times$ is the Jacobian of F . Hence, we get $\deg_{\mathbf{w}} df_1 \wedge \cdots \wedge df_n = |\mathbf{w}|$. Recall that $h_1, \dots, h_r$ are algebraically independent over k if and only if $dh_1 \wedge \cdots \wedge dh_r$ is nonzero for each $h_1, \dots, h_r \in k[\mathbf{x}]$ with $r \in \mathbf{N}$ (cf. [16, Section 26]). Thus, $f_1^{\mathbf{w}}, \dots, f_n^{\mathbf{w}}$ are algebraically independent over k if and only if $df_1^{\mathbf{w}} \wedge \cdots \wedge df_n^{\mathbf{w}}$ is nonzero, and hence if and only if

$$|\mathbf{w}| = \deg_{\mathbf{w}} df_1 \wedge \cdots \wedge df_n = \deg_{\mathbf{w}} f_1 + \cdots + \deg_{\mathbf{w}} f_n = \deg_{\mathbf{w}} F. \quad (2.2)$$

Now, assume that $n = 3$. We consider when $(d_1, d_2, d_3) \in (\Gamma_+)^3$ does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$. Note that, if d_1 and d_2 are linearly dependent over $\mathbf{Z}$, then there exists unique $(u_1, u_2) \in \mathbf{N}^2$ such that $\gcd(u_1, u_2) = 1$ and $u_2 d_1 = u_1 d_2$. Then, we can find $d \in \Gamma_+$ such that $d_i = u_i d$ for $i = 1, 2$. In this notation, we define $\gcd(d_1, d_2) = d$ and $\text{lcm}(d_1, d_2) = u_1 u_2 d$. By definition, these elements of Γ_+ are the same as the greatest common divisor and least common multiple if $\Gamma = \mathbf{Z}$. For $d, e \in \Gamma_+$, we define $\Delta_{\mathbf{w}}(d, e)$ to be the minimum among $\deg_{\mathbf{w}} df_1 \wedge df_2$ for $F \in T_3(k)$ such that $\deg_{\mathbf{w}} f_1 = d$ and $\deg_{\mathbf{w}} f_2 = e$. Here, the minimum of the empty set is defined to be ∞ . If $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, 1, 1)$, then we have $\Delta_{\mathbf{w}}(d, e) < \infty$ for any $d, e \in \mathbf{N}$, since (d, e, e) belongs to $\text{mdeg } T_3(k)$ by Proposition 1.2. For $N_1, \dots, N_r \subset \mathbf{Z}$ and $e_1, \dots, e_r \in \Gamma$ with $r \in \mathbf{N}$, we define

$$N_1 e_1 + \cdots + N_r e_r := \left\{ \sum_{i=1}^r u_i e_i \mid u_i \in N_i \text{ for } i = 1, \dots, r \right\}.$$

With the notation above, consider the following conditions:

- (K1) $d_1 < d_2 < d_3$ and $d_1 + d_2 + d_3 > |\mathbf{w}|$.
- (K2) d_2 and d_3 do not belong to $\mathbf{N}d_1$ and $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$, respectively.
- (K3) $3d_2 \neq 2d_3$ or $d_1 + d_2 < d_3 + \Delta_{\mathbf{w}}(d_2, d_3)$.

(K4) $sd_1 \neq 2d_3$ for any odd number $s \geq 3$ or $d_1 + d_2 < d_3 + \Delta_{\mathbf{w}}(d_1, d_3)$.

(K5) $4d_1 \neq 3d_2$ or $d_3 < 5 \gcd(d_1, d_2) + \Delta_{\mathbf{w}}(2d_1, d_2)$.

Then, we have the following theorem, which will be proved in Section 3 by means of the generalized Shestakov-Umirbaev theory.

Theorem 2.2. *If $d_1, d_2, d_3 \in \Gamma_+$ and $\mathbf{w} \in (\Gamma_+)^3$ satisfy (K1) through (K4), then the following assertions hold:*

(i) *Let $F \in \text{Aut}_k k[\mathbf{x}]$ be such that $\text{mdeg}_{\mathbf{w}} F = (d_1, d_2, d_3)$. If d_1 and d_2 are linearly dependent over $\mathbf{Z}$, and d_1, d_2 and d_3 satisfy (K5) and*

$$d_1 + d_2 + d_3 < \text{lcm}(d_1, d_2) + \deg_{\mathbf{w}} df_1 \wedge df_2, \quad (2.3)$$

then F does not belong to $T_3(k)$.

(ii) *If d_1 and d_2 are linearly independent over $\mathbf{Z}$, then (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$.*

As a consequence of Theorem 2.2 (i), we obtain the following theorem immediately.

Theorem 2.3. *Assume that $d_1, d_2, d_3 \in \Gamma_+$ and $\mathbf{w} \in (\Gamma_+)^3$ satisfy (K1) through (K5). If d_1 and d_2 are linearly dependent over $\mathbf{Z}$, and*

$$d_1 + d_2 + d_3 < \text{lcm}(d_1, d_2) + \Delta_{\mathbf{w}}(d_1, d_2),$$

then (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$.

Actually, if there exists $F \in T_3(k)$ such that $\text{mdeg}_{\mathbf{w}} F = (d_1, d_2, d_3)$ under the assumption of Theorem 2.3, then F satisfies (2.3), since

$$d_1 + d_2 + d_3 < \text{lcm}(d_1, d_2) + \Delta_{\mathbf{w}}(d_1, d_2) \leq \text{lcm}(d_1, d_2) + \deg_{\mathbf{w}} df_1 \wedge df_2.$$

Hence, F does not belong to $T_3(k)$ by Theorem 2.2 (i), a contradiction.

Theorem 2.2 (ii) implies the following useful criterion used in [13] to prove many interesting theorems.

Theorem 2.4 (cf. [13, Section 5]). *Let $\mathbf{w} \in (\Gamma_+)^3$ be such that w_1, w_2 and w_3 are linearly independent over $\mathbf{Z}$. If $d_1, d_2, d_3 \in \Gamma_+$ satisfy the following two conditions, then (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$:*

(1) *d_1, d_2 and d_3 are linearly dependent over $\mathbf{Z}$, and are pairwise linearly independent over $\mathbf{Z}$.*

(2) *d_i does not belong to $\mathbf{N}_0 d_j + \mathbf{N}_0 d_l$ for $(i, j, l) = (1, 2, 3), (2, 3, 1), (3, 1, 2)$.*

To check this theorem, we may assume that $(d_1, d_2, d_3) = \deg_{\mathbf{w}} F$ for some $F \in \text{Aut}_k k[\mathbf{x}]$. Then, $f_1^{\mathbf{w}}$, $f_2^{\mathbf{w}}$ and $f_3^{\mathbf{w}}$ are monomials, since w_1 , w_2 and w_3 are linearly independent over $\mathbf{Z}$. Moreover, d_1 , d_2 and d_3 are linearly dependent over $\mathbf{Z}$ if and only if $f_1^{\mathbf{w}}$, $f_2^{\mathbf{w}}$ and $f_3^{\mathbf{w}}$ are algebraically dependent over k , and hence if and only if $d_1 + d_2 + d_3 = \deg_{\mathbf{w}} F > |\mathbf{w}|$ by (2.2) and the preceding discussion. Using this remark, we can easily check that $d_{\sigma(1)}$, $d_{\sigma(2)}$ and $d_{\sigma(3)}$ satisfy (K1) through (K4) for some $\sigma \in \mathfrak{S}_3$ if (1) and (2) are satisfied.

Since $\Delta_{\mathbf{w}}(d, e)$'s are not easy to estimate in general, Theorems 2.2 and 2.3 are not suitable for practical use. So we derive a theorem which is useful in applications. Recall that $\mathbf{N}_0 e_1 + \cdots + \mathbf{N}_0 e_r$ is a well-ordered subset of Γ for each $e_1, \dots, e_r \in \Gamma_+$ with $r \in \mathbf{N}$ (cf. [12, Lemma 6.1 (ii)]). Hence, we can define

$$|\mathbf{w}|_* := \min(\{\gamma \in \mathbf{N}w_{\sigma(1)} + \mathbf{N}w_{\sigma(2)} \mid \gamma > w_{\sigma(1)} + w_{\sigma(3)}\} \cup \{2w_{\sigma(1)} + w_{\sigma(3)}\}),$$

where $\sigma \in \mathfrak{S}_3$ is such that $w_{\sigma(1)} \leq w_{\sigma(2)} \leq w_{\sigma(3)}$.

Consider the following conditions:

(A) One of the following holds:

(A1) $3d_2 \neq 2d_3$ and $sd_1 \neq 2d_3$ for any odd number $s \geq 3$.

(A2) $3d_2 = 2d_3$ and $d_1 + d_2 < d_3 + \max\{\Delta_{\mathbf{w}}(d_2, d_3), |\mathbf{w}|_*\}$.

(A3) $sd_1 = 2d_3$ for some odd number $s \geq 3$ and

$$d_1 + d_2 < d_3 + \max\{\Delta_{\mathbf{w}}(d_1, d_3), |\mathbf{w}|_*\}.$$

(B) d_1 and d_2 are linearly independent over $\mathbf{Z}$, or d_1 and d_2 are linearly dependent over $\mathbf{Z}$ and one of the following holds:

(B1) $\gcd(d_1, d_2) \leq |\mathbf{w}|_*$ and d_3 belongs to $\mathbf{N} \gcd(d_1, d_2)$.

(B2) $d_1 + d_2 + d_3 < \text{lcm}(d_1, d_2) + |\mathbf{w}|_*$.

The following is our main theorem.

Theorem 2.5. *If $d_1, d_2, d_3 \in \Gamma_+$ and $\mathbf{w} \in (\Gamma_+)^3$ satisfy (K1), (K2), (A) and (B), then (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$.*

We prove this theorem in Section 5 using Theorem 2.2 and a weighted version of Sun-Chen's lemma (Theorem 4.1).

To end this section, we show Theorem 1.3 by assuming Theorem 2.5. Take any $d_1, d_2, d_3 \in \mathbf{N}$ with $d_1 \leq d_2 \leq d_3$ which satisfy (a), (b) and (c). It suffices to check that (K1), (K2), (A) and (B) hold for $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, 1, 1)$. Since $d_1 \leq d_2 \leq d_3$, we have $d_1 < d_2 < d_3$ by (c). Since $d_1 \geq 1$, it follows that $d_1 + d_2 + d_3 > 3 = |\mathbf{w}|$, proving (K1). Clearly, (c) implies (K2). If (a1)

holds, then we have (A1). Assume that (a1) does not hold. Then, we have $3d_2 = 2d_3$ or $sd_1 = 2d_3$ for some odd number $s \geq 3$, and (a2) is satisfied. Since $|\mathbf{w}|_* = 3$, it follows that

$$d_1 + d_2 \leq d_3 + 2 < d_3 + |\mathbf{w}|_*.$$

Thus, we get (A2) or (A3). Since $|\mathbf{w}|_* = 3$, (b1) and (b2) imply (B1) and (B2), respectively. Therefore, d_1 , d_2 and d_3 satisfy (K1), (K2), (A) and (B).

3 Generalized Shestakov-Umirbaev theory

The goal of this section is to prove Theorem 2.2. First, we briefly recall the generalized Shestakov-Umirbaev theory [11], [12]. For the moment, $n \in \mathbf{N}$ may be arbitrary. Let $\mathbf{w}$ be an element of $(\Gamma_+)^n$, and let $F = (f_1, f_2, f_3)$ and $G = (g_1, g_2, g_3)$ be triples of elements of $k[\mathbf{x}]$ such that f_1, f_2, f_3 and g_1, g_2, g_3 are algebraically independent over k , respectively. We say that the pair (F, G) satisfies the *Shestakov-Umirbaev condition* for the weight $\mathbf{w}$ if the following conditions hold (cf. [12]):

- (SU1) $g_1 = f_1 + af_3^2 + cf_3$ and $g_2 = f_2 + bf_3$ for some $a, b, c \in k$, and $g_3 - f_3$ belongs to $k[g_1, g_2]$;
- (SU2) $\deg_{\mathbf{w}} f_1 \leq \deg_{\mathbf{w}} g_1$ and $\deg_{\mathbf{w}} f_2 = \deg_{\mathbf{w}} g_2$;
- (SU3) $(g_1^{\mathbf{w}})^2 \approx (g_2^{\mathbf{w}})^s$ for some odd number $s \geq 3$;
- (SU4) $\deg_{\mathbf{w}} f_3 \leq \deg_{\mathbf{w}} g_1$, and $f_3^{\mathbf{w}}$ does not belong to $k[g_1^{\mathbf{w}}, g_2^{\mathbf{w}}]$;
- (SU5) $\deg_{\mathbf{w}} g_3 < \deg_{\mathbf{w}} f_3$;
- (SU6) $\deg_{\mathbf{w}} g_3 < \deg_{\mathbf{w}} g_1 - \deg_{\mathbf{w}} g_2 + \deg_{\mathbf{w}} dg_1 \wedge dg_2$.

Here, $h_1 \approx h_2$ (resp. $h_1 \not\approx h_2$) denotes that h_1 and h_2 are linearly dependent (resp. linearly independent) over k for each $h_1, h_2 \in k[\mathbf{x}] \setminus \{0\}$. We say that (F, G) satisfies the *weak Shestakov-Umirbaev condition* for the weight $\mathbf{w}$ if (SU4), (SU5), (SU6) and the following conditions are satisfied (cf. [12]):

- (SU1') $g_1 - f_1$, $g_2 - f_2$ and $g_3 - f_3$ belong to $k[f_2, f_3]$, $k[f_3]$ and $k[g_1, g_2]$, respectively;
- (SU2') $\deg f_i \leq \deg g_i$ for $i = 1, 2$;
- (SU3') $\deg g_2 < \deg g_1$, and $g_1^{\mathbf{w}}$ does not belong to $k[g_2^{\mathbf{w}}]$.

It is easy to check that (SU1), (SU2) and (SU3) imply (SU1'), (SU2') and (SU3'), respectively. Hence, the Shestakov-Umirbaev condition implies the weak Shestakov-Umirbaev condition. If (F, G) satisfies the weak Shestakov-Umirbaev condition for the weight $\mathbf{w}$, then (F, G) has the following properties due to [12, Theorem 4.2]. Here, we regard Γ as a subgroup of $\mathbf{Q} \otimes_{\mathbf{Z}} \Gamma$ which

has a structure of totally ordered additive group induced from Γ . We note that δ in (P1) is equal to $\gcd(\deg_{\mathbf{w}} g_1, \deg_{\mathbf{w}} g_2)$ in our notation.

(P1) $(g_1^{\mathbf{w}})^2 \approx (g_2^{\mathbf{w}})^s$ for some odd number $s \geq 3$, and so $\delta := (1/2) \deg_{\mathbf{w}} g_2$ belongs to Γ .

(P2) $\deg_{\mathbf{w}} f_3 \geq (s-2)\delta + \deg_{\mathbf{w}} dg_1 \wedge dg_2$.

(P5) If $\deg_{\mathbf{w}} f_1 < \deg_{\mathbf{w}} g_1$, then $s = 3$, $g_1^{\mathbf{w}} \approx (f_3^{\mathbf{w}})^2$, $\deg_{\mathbf{w}} f_3 = (3/2)\delta$ and

$$\deg_{\mathbf{w}} f_1 \geq \frac{5}{2}\delta + \deg_{\mathbf{w}} dg_1 \wedge dg_2.$$

(P7) $\deg_{\mathbf{w}} f_2 < \deg_{\mathbf{w}} f_1$, $\deg_{\mathbf{w}} f_3 \leq \deg_{\mathbf{w}} f_1$, and $\delta < \deg_{\mathbf{w}} f_i \leq s\delta$ for $i = 1, 2, 3$.

(P11) If $\deg_{\mathbf{w}} f_3 \leq \deg_{\mathbf{w}} f_2$, then $s = 3$.

We say that $F \in \text{Aut}_k k[\mathbf{x}]$ admits an *elementary reduction* for the weight $\mathbf{w}$ if there exists an elementary automorphism E such that $\deg_{\mathbf{w}} F \circ E < \deg_{\mathbf{w}} F$. For a k -vector subspace V of $k[\mathbf{x}]$, we define $V^{\mathbf{w}}$ to be the k -vector subspace of $k[\mathbf{x}]$ generated by $\{f^{\mathbf{w}} \mid f \in V\}$. Then, $F \in \text{Aut}_k k[\mathbf{x}]$ admits an elementary reduction for the weight $\mathbf{w}$ if and only if $f_l^{\mathbf{w}}$ belongs to $k[\{f_i \mid i \neq l\}]^{\mathbf{w}}$ for some $1 \leq l \leq n$.

Now, assume that $n = 3$. For $\sigma \in \mathfrak{S}_3$ and $F \in \text{Aut}_k k[\mathbf{x}]$, we define $F_{\sigma} = (f_{\sigma(1)}, f_{\sigma(2)}, f_{\sigma(3)})$. We say that F admits a *Shestakov-Umirbaev reduction* for the weight $\mathbf{w}$ if there exist $G \in \text{Aut}_k k[\mathbf{x}]$ and $\sigma \in \mathfrak{S}_3$ such that (F_{σ}, G_{σ}) satisfies the Shestakov-Umirbaev condition for the weight $\mathbf{w}$.

The following theorem is a generalization of the main result of Shestakov-Umirbaev [19].

Theorem 3.1 ([12, Theorem 2.1]). *If $\deg_{\mathbf{w}} F > |\mathbf{w}|$ holds for $F \in T_3(k)$ and $\mathbf{w} \in (\Gamma_+)^3$, then F admits an elementary reduction or a Shestakov-Umirbaev reduction for the weight $\mathbf{w}$.*

We also use the following version of the Shestakov-Umirbaev inequality (see [12, Section 3] for detail). Let $S = \{f, g\}$ be a subset of $k[\mathbf{x}]$ such that f and g are algebraically independent over k , and h a nonzero element of $k[S]$. Then, we can uniquely express $h = \sum_{i,j} c_{i,j} f^i g^j$, where $c_{i,j} \in k$ for each $i, j \in \mathbf{N}_0$. We define $\deg_{\mathbf{w}}^S h$ to be the maximum among $\deg_{\mathbf{w}} f^i g^j$ for $i, j \in \mathbf{N}_0$ with $c_{i,j} \neq 0$. We note that, if $h^{\mathbf{w}}$ does not belong to $k[f^{\mathbf{w}}, g^{\mathbf{w}}]$, or if $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}} g$ and $h^{\mathbf{w}}$ does not belong to $k[f^{\mathbf{w}}]$, then $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}}^S h$.

With the notation and assumption above, the following lemma holds (see [12, Lemmas 3.2 (i) and 3.3 (ii)] for the proof).

Lemma 3.2. *If $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}}^S h$, then there exist $p, q \in \mathbf{N}$ with $\gcd(p, q) = 1$ such that $(g^{\mathbf{w}})^p \approx (f^{\mathbf{w}})^q$ and*

$$\deg_{\mathbf{w}} h \geq q \deg_{\mathbf{w}} f - \deg_{\mathbf{w}} f - \deg_{\mathbf{w}} g + \deg_{\mathbf{w}} df \wedge dg.$$

Assume further that $\deg_{\mathbf{w}} f < \deg_{\mathbf{w}} g$ and $g^{\mathbf{w}}$ does not belong to $k[f^{\mathbf{w}}]$. If $\deg_{\mathbf{w}} h \leq \deg_{\mathbf{w}} g$, then we have $p = 2$, and q is an odd number with $q \geq 3$. If $\deg_{\mathbf{w}} h \leq \deg_{\mathbf{w}} f$, then we have $q = 3$.

Now, we prove Theorem 2.2. First, consider the following conditions for $e_1, e_2, e_3 \in \Gamma_+$ and $\mathbf{w} \in (\Gamma_+)^3$:

(K2') e_1 and e_2 do not belong to $\mathbf{N}_0 e_2 + \mathbf{N}_0 e_3$ and $\mathbf{N} e_3$, respectively.

(K3') $2e_1 = 3e_2$ and $e_2 + e_3 \geq e_1 + \Delta_{\mathbf{w}}(e_1, e_2)$.

(K4') $2e_1 = se_3$ for some odd number $s \geq 3$ and $e_2 + e_3 \geq e_1 + \Delta_{\mathbf{w}}(e_1, e_3)$.

(K5') $3e_2 = 4e_3$ and $e_1 \geq 5 \gcd(e_2, e_3) + \Delta_{\mathbf{w}}(e_2, 2e_3)$.

The following lemma is a direct consequence of the generalized Shestakov-Umirbaev theory.

Lemma 3.3. *Let $e_1, e_2, e_3 \in \Gamma_+$ and $\mathbf{w} \in (\Gamma_+)^3$ be such that $e_1 > e_2 > e_3$ and $\text{mdeg}_{\mathbf{w}} F = (e_1, e_2, e_3)$ for some $F \in T_3(k)$.*

(i) *If F admits a Shestakov-Umirbaev reduction for the weight $\mathbf{w}$, then one of (K3'), (K4') and (K5') holds.*

(ii) *Assume that (K2') is satisfied. Then, the following assertions hold:*

(1) *If $f_3^{\mathbf{w}}$ belongs to $k[f_1, f_2]^{\mathbf{w}}$, then we have (K3').*

(2) *If $f_2^{\mathbf{w}}$ belongs to $k[f_1, f_3]^{\mathbf{w}}$, then we have (K4').*

(3) *If $f_1^{\mathbf{w}}$ belongs to $k[f_2, f_3]^{\mathbf{w}}$, then e_2 and e_3 are linearly dependent over $\mathbf{Z}$ and*

$$e_1 + e_2 + e_3 \geq \text{lcm}(e_2, e_3) + \deg_{\mathbf{w}} df_2 \wedge df_3. \quad (3.1)$$

Proof. (i) By assumption, there exist $G \in \text{Aut}_k k[\mathbf{x}]$ and $\sigma \in \mathfrak{S}_3$ such that (F_{σ}, G_{σ}) satisfies the Shestakov-Umirbaev condition for the weight $\mathbf{w}$. Then, we see from (SU1) that G is tame, since so is F . Since $e_1 > e_i$ for $i = 2, 3$, we have $\sigma(1) = 1$ in view of (P7). Hence, σ must be the identity permutation id or the transposition $(2, 3)$. First, assume that $\sigma = \text{id}$. Then, we have $s = 3$ by (P11), since $e_2 > e_3$. By (SU2), we get $\deg_{\mathbf{w}} g_1 \leq e_1$ and $\deg_{\mathbf{w}} g_2 = e_2$. When $\deg_{\mathbf{w}} g_1 = e_1$, (SU3) implies

$$2e_1 = 2 \deg_{\mathbf{w}} g_1 = s \deg_{\mathbf{w}} g_2 = 3 \deg_{\mathbf{w}} g_2 = 3e_2.$$

Since G is tame, and $\deg_{\mathbf{w}} g_i = e_i$ for $i = 1, 2$, we know by (P2) that

$$e_3 \geq (3 - 2) \frac{1}{2} e_2 + \deg_{\mathbf{w}} dg_1 \wedge dg_2 \geq e_1 - e_2 + \Delta_{\mathbf{w}}(e_1, e_2).$$

Therefore, (K3') holds in this case. If $\deg_{\mathbf{w}} g_1 < e_1$, then we have $\deg_{\mathbf{w}} g_1 = 2e_3$ and $e_3 = (3/2)(1/2)e_2$ by (P5). Hence, we get $3e_2 = 4e_3$, $\gcd(e_2, e_3) = (1/4)e_2$ and $\deg_{\mathbf{w}} dg_1 \wedge dg_2 \geq \Delta_{\mathbf{w}}(2e_3, e_2)$. Thus, the last part of (P5) implies

$$e_1 \geq \frac{5}{2} \frac{1}{2} e_2 + \deg_{\mathbf{w}} dg_1 \wedge dg_2 \geq 5 \gcd(e_2, e_3) + \Delta_{\mathbf{w}}(2e_3, e_2).$$

Therefore, (K5') holds in this case. Next, assume that $\sigma = (2, 3)$. Then, we have $\deg_{\mathbf{w}} g_3 = \deg_{\mathbf{w}} f_3 = e_3$ by (SU2). We also have $\deg_{\mathbf{w}} g_1 = \deg_{\mathbf{w}} f_1 = e_1$, for otherwise $e_2 = (3/2)(1/2)e_3$ by (P5), and so $e_2 < e_3$, a contradiction. Hence, it follows from (SU3) that $2e_1 = se_3$ for some odd number $s \geq 3$. Thus, we get

$$e_2 \geq (s-2) \frac{1}{2} e_3 + \deg_{\mathbf{w}} dg_1 \wedge dg_3 \geq e_1 - e_3 + \Delta_{\mathbf{w}}(e_1, e_3)$$

by (P2). Therefore, (K4') holds in this case.

(ii) In the case of (1), there exists $h \in k[f_1, f_2]$ such that $h^{\mathbf{w}} = f_3^{\mathbf{w}}$. Then, we have $\deg_{\mathbf{w}} h = e_3 < e_i$ for $i = 1, 2$. Since $h^{\mathbf{w}} = f_3^{\mathbf{w}}$ is not a constant, it follows that $h^{\mathbf{w}}$ does not belong to $k[f_1^{\mathbf{w}}, f_2^{\mathbf{w}}]$. This implies that $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}}^S h$ holds for $S = \{f_1, f_2\}$ as mentioned. By Lemma 3.2, there exist $p, q \in \mathbb{N}$ with $\gcd(p, q) = 1$ such that $(f_1^{\mathbf{w}})^p \approx (f_2^{\mathbf{w}})^q$ and

$$\begin{aligned} e_3 = \deg_{\mathbf{w}} h &\geq q \deg_{\mathbf{w}} f_2 - \deg_{\mathbf{w}} f_1 - \deg_{\mathbf{w}} f_2 + \deg_{\mathbf{w}} df_1 \wedge df_2 \\ &\geq qe_2 - e_1 - e_2 + \Delta_{\mathbf{w}}(e_1, e_2). \end{aligned} \quad (3.2)$$

We note that $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}} f_2 < \deg_{\mathbf{w}} f_1$, and $f_1^{\mathbf{w}}$ does not belong to $k[f_2^{\mathbf{w}}]$ by (K2'). Hence, we have $(p, q) = (2, 3)$ by the last part of Lemma 3.2. Since $(f_1^{\mathbf{w}})^p \approx (f_2^{\mathbf{w}})^q$, it follows that $2e_1 = 3e_2$, and thus $qe_2 - e_1 - e_2 = e_1 - e_2$. Therefore, we obtain (K3') from (3.2).

We prove (2) similarly. Let $h \in k[f_1, f_3]$ be such that $h^{\mathbf{w}} = f_2^{\mathbf{w}}$. Then, we have $\deg_{\mathbf{w}} h = e_2 < e_1 = \deg_{\mathbf{w}} f_1$. Since e_2 does not belong to $\mathbb{N}e_3$ by (K2'), $h^{\mathbf{w}}$ does not belong to $k[f_3^{\mathbf{w}}]$. Hence, $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}}^S h$ holds for $S = \{f_1, f_3\}$ as mentioned. As before, there exist $p, q \in \mathbb{N}$ with $\gcd(p, q) = 1$ such that $pe_1 = qe_3$ and

$$e_2 = \deg_{\mathbf{w}} h \geq qe_3 - e_1 - e_3 + \Delta_{\mathbf{w}}(e_1, e_3) \quad (3.3)$$

by Lemma 3.2. We note that $\deg_{\mathbf{w}} f_3$ and $\deg_{\mathbf{w}} h$ are less than $\deg_{\mathbf{w}} f_1$, and $f_1^{\mathbf{w}}$ does not belong to $k[f_3^{\mathbf{w}}]$ by (K2'). Hence, we know that $p = 2$, and q is an odd number with $q \geq 3$ by the last part of Lemma 3.2. Thus, we get $2e_1 = qe_3$, and so $qe_3 - e_1 - e_3 = e_1 - e_3$. Therefore, we obtain (K4') from (3.3).

Finally, we prove (3). Let $h \in k[f_2, f_3]$ be such that $h^{\mathbf{w}} = f_1^{\mathbf{w}}$. Then, we have $\deg_{\mathbf{w}} h = e_1$. Since e_1 does not belong to $\mathbf{N}_0 e_2 + \mathbf{N}_0 e_3$ by (K2'), it follows that $h^{\mathbf{w}}$ does not belong to $k[f_2^{\mathbf{w}}, f_3^{\mathbf{w}}]$. This implies that $\deg_{\mathbf{w}} h < \deg_{\mathbf{w}}^S h$ holds for $S = \{f_2, f_3\}$. As before, there exist $p, q \in \mathbf{N}$ with $\gcd(p, q) = 1$ such that $pe_3 = qe_2$ and $e_1 = \deg_{\mathbf{w}} h \geq qe_2 - e_2 - e_3 + \deg_{\mathbf{w}} df_2 \wedge df_3$. Since $qe_2 = \text{lcm}(e_2, e_3)$, this inequality yields (3.1), proving (3). $\square$

Now, let us complete the proof of Theorem 2.2. Set $e_1 := d_3$, $e_2 := d_2$ and $e_3 := d_1$. Then, we have $e_1 > e_2 > e_3$ by (K1). First, we prove (i) by contradiction. Suppose that F is tame. Then, $F' := (f_3, f_2, f_1)$ is also tame. Since $\deg_{\mathbf{w}} F' > |\mathbf{w}|$ by (K1), it follows that F' admits a Shestakov-Umirbaev reduction or an elementary reduction for the weight $\mathbf{w}$ by Theorem 3.1. Since $\text{mdeg}_{\mathbf{w}} F' = (e_1, e_2, e_3)$, we know by Lemma 3.3 that one of (K3'), (K4'), (K5') and (3.1) must be satisfied. This contradicts (K3), (K4), (K5) and (2.3). Therefore, F does not belong to $T_3(k)$, proving (i).

We prove (ii) similarly. Suppose that (d_1, d_2, d_3) belongs to $\text{mdeg}_{\mathbf{w}} T_3(k)$. Then, (e_1, e_2, e_3) also belongs to $\text{mdeg}_{\mathbf{w}} T_3(k)$. Hence, there exists $F' \in T_3(k)$ such that $\text{mdeg}_{\mathbf{w}} F' = (e_1, e_2, e_3)$. As in the case of (i), this implies that one of (K3'), (K4') and (K5') holds, or e_2 and e_3 are linearly dependent over $\mathbf{Z}$ by Lemma 3.3. However, (K3') and (K4') do not hold because of (K3) and (K4). Since $e_2 = d_2$ and $e_3 = d_1$ are linearly independent over $\mathbf{Z}$ by assumption, the rest of the conditions also do not hold. This is a contradiction. Therefore, (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$, proving (ii). This completes the proof of Theorem 2.2.

4 A generalization of Sun-Chen's lemma

In this section, we discuss lower bounds for the degrees of differential forms. Note that, if $df_1 \wedge \cdots \wedge df_r \neq 0$ for $f_1, \dots, f_r \in k[\mathbf{x}]$ with $r \geq 1$, then we have $\deg df_1 \wedge \cdots \wedge df_r \geq r$ by definition. In the case of $r = 2$, Karaš [8, Theorem 14] showed that $\deg df_1 \wedge df_2 \geq 4$ under the additional assumption that $\deg f_1 = 4$ and $\deg f_2 = 6$. When $n = 3$, Sun-Chen [20, Lemma 3.1] showed that $\deg df_1 \wedge df_2 \geq 3$ for each $F \in \text{Aut}_k k[\mathbf{x}]$ with $\deg f_1 \nmid \deg f_2$ and $\deg f_2 \nmid \deg f_1$.

The purpose of this section is to prove the following theorem. This theorem is considered as a generalization of Sun-Chen's lemma, since $|\mathbf{w}|_* = 3$ if $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, 1, 1)$.

Theorem 4.1. *Assume that $n = 3$. Let $F \in \text{Aut}_k k[\mathbf{x}]$ and $\mathbf{w} \in (\Gamma_+)^3$ be such that $\deg_{\mathbf{w}} f_1$ or $\deg_{\mathbf{w}} f_2$ does not belong to $\{w_1, w_2, w_3\}$, and $\deg_{\mathbf{w}} f_i$*

does not belong to $\mathbf{N} \deg_{\mathbf{w}} f_j$ for each $(i, j) \in \{(1, 2), (2, 1)\}$. Then, we have $\deg_{\mathbf{w}} df_1 \wedge df_2 \geq |\mathbf{w}|_*$.

As a consequence of Theorem 4.1, we know that $\Delta_{\mathbf{w}}(d_1, d_2) \geq |\mathbf{w}|_*$ holds for $d_1, d_2 \in \Gamma_+$ and $\mathbf{w} \in (\Gamma_+)^3$ if d_1 or d_2 does not belong to $\{w_1, w_2, w_3\}$, and d_i does not belong to $\mathbf{N}d_j$ for each $(i, j) \in \{(1, 2), (2, 1)\}$. We also note that Karaš [8, Theorem 14] implies $\Delta_{\mathbf{w}}(4, 6) \geq 4$ when $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, 1, 1)$.

To prove Theorem 4.1, we need the following two propositions.

Proposition 4.2. *Assume that $n = 2$. Let $F \in \text{Aut}_k k[\mathbf{x}]$ and $\mathbf{w} \in (\Gamma_+)^2$ be such that $\deg_{\mathbf{w}} F > |\mathbf{w}|$. Then, there exist $(i, j) \in \{(1, 2), (2, 1)\}$ and $l \in \mathbf{N}$ such that $f_i^{\mathbf{w}} \approx (f_j^{\mathbf{w}})^l$. Hence, $\deg_{\mathbf{w}} f_i$ belongs to $\mathbf{N} \deg_{\mathbf{w}} f_j$.*

Proof. Since $\deg_{\mathbf{w}} F > |\mathbf{w}|$ by assumption, $f_1^{\mathbf{w}}$ and $f_2^{\mathbf{w}}$ are algebraically dependent over k by (2.2) and the preceding discussion. Hence, x_l does not belong to $k[f_1^{\mathbf{w}}, f_2^{\mathbf{w}}]$ for some $l \in \{1, 2\}$. On the other hand, x_l belongs to $k[\mathbf{x}] = k[f_1, f_2]$. Thus, $\deg_{\mathbf{w}}^S x_l > \deg_{\mathbf{w}} x_l$ holds for $S = \{f_1, f_2\}$. By Lemma 3.2, there exist $p_1, p_2 \in \mathbf{N}$ with $\gcd(p_1, p_2) = 1$ such that $(f_2^{\mathbf{w}})^{p_1} \approx (f_1^{\mathbf{w}})^{p_2}$ and

$$w_l = \deg_{\mathbf{w}} x_l \geq p_2 d_1 - d_1 - d_2 + \deg_{\mathbf{w}} df_1 \wedge df_2,$$

where $d_i := \deg_{\mathbf{w}} f_i$ for $i = 1, 2$. Then, we have $d_i = p_i \gcd(d_1, d_2)$ for $i = 1, 2$. Since $\deg_{\mathbf{w}} df_1 \wedge df_2 = |\mathbf{w}|$, it follows that

$$(p_1 + p_2 - p_1 p_2) \gcd(d_1, d_2) = d_1 + d_2 - p_2 d_1 \geq |\mathbf{w}| - w_l \geq \min\{w_1, w_2\} > 0.$$

Hence, we get $p_1 p_2 < p_1 + p_2$. Since $\gcd(p_1, p_2) = 1$, this implies that $p_1 = 1$ or $p_2 = 1$. Therefore, we have $f_2^{\mathbf{w}} \approx (f_1^{\mathbf{w}})^{p_2}$ or $(f_2^{\mathbf{w}})^{p_1} \approx f_1^{\mathbf{w}}$. $\square$

Next, assume that $n \geq 3$, and let $\mathbf{w} \in (\Gamma_+)^n$ be such that $w_1 \leq \dots \leq w_n$. We define

$$|\mathbf{w}|_* := \min(\{\gamma \in \mathbf{N}w_1 + \dots + \mathbf{N}w_{n-1} \mid \gamma > |\mathbf{w}| - w_{n-1}\} \cup \{|\mathbf{w}| + w_1 - w_{n-1}\}).$$

Note that, if $n = 3$, then this is the same as $|\mathbf{w}|_*$ defined before.

With this notation, we have the following proposition.

Proposition 4.3. *If $f_1, \dots, f_{n-1} \in k[\mathbf{x}]$ are algebraically independent over k and $\deg_{\mathbf{w}} df_1 \wedge \dots \wedge df_{n-1} < |\mathbf{w}|_*$, then there exist $\phi \in \mathbf{T}_n(k)$ and $1 \leq l \leq n$ such that $\phi(f_i)$ belongs to $k[\mathbf{x} \setminus \{x_l\}]$ and $\deg_{\mathbf{w}} \phi(f_i) = \deg_{\mathbf{w}} f_i$ for $i = 1, \dots, n-1$.*

Before proving Proposition 4.3, we show how Propositions 4.2 and 4.3 imply Theorem 4.1. By changing the indices of x_1, x_2 and x_3 , and w_1, w_2 and w_3 if necessary, we may assume that $w_1 \leq w_2 \leq w_3$. Suppose to the contrary that $\deg_{\mathbf{w}} df_1 \wedge df_2 < |\mathbf{w}|_*$ for some $F \in \text{Aut}_k k[\mathbf{x}]$ and $\mathbf{w} \in (\Gamma_+)^3$ which satisfy the assumptions of Theorem 4.1. By Proposition 4.3, there exist $\phi \in T_3(k)$ and $1 \leq i_1 < i_2 \leq 3$ such that $\phi(f_i)$ belongs to $k[x_{i_1}, x_{i_2}]$ and $\deg_{\mathbf{w}} \phi(f_i) = \deg_{\mathbf{w}} f_i$ for $i = 1, 2$. Then, $G := (\phi(f_1), \phi(f_2))$ belongs to $\text{Aut}_k k[x_{i_1}, x_{i_2}]$. Set $\mathbf{v} := (w_{i_1}, w_{i_2})$. Then, we have

$$d_i := \deg_{\mathbf{w}} f_i = \deg_{\mathbf{w}} \phi(f_i) = \deg_{\mathbf{v}} \phi(f_i)$$

for $i = 1, 2$. If $\deg_{\mathbf{v}} G = |\mathbf{v}|$, then $(d_1, d_2) = \text{mdeg}_{\mathbf{v}} G$ is equal to (w_{i_1}, w_{i_2}) or (w_{i_2}, w_{i_1}) by the remark after Lemma 2.1. Hence, d_1 and d_2 both belong to $\{w_1, w_2, w_3\}$, a contradiction. If $\deg_{\mathbf{v}} G > |\mathbf{v}|$, then d_i belongs to $\mathbf{N}d_j$ for some $(i, j) \in \{(1, 2), (2, 1)\}$ by Proposition 4.2, a contradiction. Therefore, Theorem 4.1 holds true.

To prove Proposition 4.3, we need some lemmas. Let $k(\mathbf{x})$ be the field of fractions of $k[\mathbf{x}]$. Then, the following assertion holds.

Lemma 4.4. *Assume that $g_1, \dots, g_r \in k(\mathbf{x})$ and $1 \leq l \leq n$ satisfy the following conditions, where $1 \leq r < n$. Then, $g_1, \dots, g_r$ belong to $k(\mathbf{x} \setminus \{x_l\})$.*

- (1) $g_1, \dots, g_r$ are algebraically independent over k .
- (2) $g_1, \dots, g_r, x_{i_1}, \dots, x_{i_{n-r}}$ are algebraically dependent over k for any distinct $i_1, \dots, i_{n-r} \in \{1, \dots, n\} \setminus \{l\}$.

Proof. Set $S = \{x_i \mid i \neq l\} \cup \{g_1, \dots, g_r\}$. If the assertion is false, then $k(\mathbf{x})$ is algebraic over $k(S)$. By (1), there exists a transcendence basis S_0 of $k(\mathbf{x})$ over k such that $\{g_1, \dots, g_r\} \subset S_0 \subset S$. Then, we have

$$S_0 = \{g_1, \dots, g_r, x_{i_1}, \dots, x_{i_{n-r}}\}$$

for some distinct $i_1, \dots, i_{n-r} \in \{1, \dots, n\} \setminus \{l\}$. This contradicts (2). $\square$

Now, write

$$\omega := df_1 \wedge \dots \wedge df_{n-1} = \sum_{i=1}^n h_i \eta_i,$$

where $h_i \in k[\mathbf{x}]$ and $\eta_i := dx_1 \wedge \dots \wedge dx_{i-1} \wedge dx_{i+1} \wedge \dots \wedge dx_n$ for each i . Then, we have $\deg_{\mathbf{w}} \omega \geq \deg_{\mathbf{w}} h_i \eta_i$ for $i = 1, \dots, n$. For each $\phi \in \text{Aut}_k k[\mathbf{x}]$, it holds that

$$\omega^\phi := d\phi(f_1) \wedge \dots \wedge d\phi(f_{n-1}) = \sum_{i=1}^n \phi(h_i) \eta_i^\phi, \quad (4.1)$$

where $\eta_i^\phi := d\phi(x_1) \wedge \cdots \wedge d\phi(x_{i-1}) \wedge d\phi(x_{i+1}) \wedge \cdots \wedge d\phi(x_n)$ for each i . In fact, we have

$$\begin{aligned} d\phi(f_i) &= d(f_i(\phi(x_1), \dots, \phi(x_n))) \\ &= \sum_{j=1}^n \left(\frac{\partial f_i}{\partial x_j} \right) (\phi(x_1), \dots, \phi(x_n)) d\phi(x_j) = \sum_{j=1}^n \phi \left(\frac{\partial f_i}{\partial x_j} \right) d\phi(x_j) \end{aligned}$$

for each i by chain rule. We note that, if ϕ is an elementary automorphism such that $\phi(x_l) = x_l + p$ for some $1 \leq l \leq n$ and $p \in k[\mathbf{x} \setminus \{x_l\}]$, then

$$\eta_l^\phi = \eta_l \quad \text{and} \quad \eta_i^\phi = \eta_i - (-1)^{i-l} \frac{\partial p}{\partial x_i} \eta_l \quad \text{for each } i \neq l. \quad (4.2)$$

With the notation above, we have the following lemma.

Lemma 4.5. *If $\deg_{\mathbf{w}} \omega < |\mathbf{w}|_*$, then there exist $\phi \in T_n(k)$, $h \in k[\mathbf{x}]$ and $1 \leq l \leq n$ such that $\omega^\phi = h\eta_l$ and $\deg_{\mathbf{w}} \phi(f_i) = \deg_{\mathbf{w}} f_i$ for $i = 1, \dots, n-1$.*

Before proving Lemma 4.5, we show Proposition 4.3 by assuming this lemma. Since $\deg_{\mathbf{w}} \omega < |\mathbf{w}|_*$ by assumption, there exist $\phi \in T_n(k)$, $h \in k[\mathbf{x}]$ and $1 \leq l \leq n$ as in Lemma 4.5. We have only to check that $\phi(f_i)$ belongs to $k[\mathbf{x} \setminus \{x_l\}]$ for $i = 1, \dots, n-1$. Since $\omega^\phi = h\eta_l$, we have

$$d\phi(f_1) \wedge \cdots \wedge d\phi(f_{n-1}) \wedge dx_i = \omega^\phi \wedge dx_i = h\eta_l \wedge dx_i = 0$$

for each $i \neq l$. Hence, $\phi(f_1), \dots, \phi(f_{n-1}), x_i$ are algebraically dependent over k for each $i \neq l$. Thus, $\phi(f_1), \dots, \phi(f_{n-1})$ belong to $k[\mathbf{x} \setminus \{x_l\}]$ by Lemma 4.4. Therefore, Proposition 4.3 follows from Lemma 4.5.

To prove Lemma 4.5, we use the following two lemmas.

Lemma 4.6. *Assume that $\sigma \in \text{Aut}_k k[\mathbf{x}]$ is such that $\sigma(x_i) = \alpha_i x_i + p_i$ for some $\alpha_i \in k^\times$ and $p_i \in k[x_1, \dots, x_{i-1}]$ with $\deg_{\mathbf{w}} p_i \leq w_i$ for $i = 1, \dots, n$. Then, $\deg_{\mathbf{w}} \sigma(f) = \deg_{\mathbf{w}} f$ holds for each $f \in k[\mathbf{x}]$.*

Proof. We may assume that $f \neq 0$. Since $\deg_{\mathbf{w}} p_i \leq w_i$ by assumption, we have $\deg_{\mathbf{w}} \sigma(x_i) = w_i$ for each i . Hence, the assertion holds when f is a monomial in view of (2.1). This implies that $\deg_{\mathbf{w}} \sigma(f) \leq \deg_{\mathbf{w}} f$ in the general case. Since $\sigma(f) = \sigma(f^{\mathbf{w}}) + \sigma(f - f^{\mathbf{w}})$ with

$$\deg_{\mathbf{w}} \sigma(f - f^{\mathbf{w}}) \leq \deg_{\mathbf{w}} (f - f^{\mathbf{w}}) < \deg_{\mathbf{w}} f,$$

it suffices to show that $\deg_{\mathbf{w}} \sigma(f^{\mathbf{w}}) = \deg_{\mathbf{w}} f$. Since $\deg_{\mathbf{w}} p_i \leq w_i$, each $\sigma(x_i)^{\mathbf{w}}$ has the form $\alpha_i x_i + p'_i$ for some $p'_i \in k[x_1, \dots, x_{i-1}]$. Define $\tau \in \text{Aut}_k k[\mathbf{x}]$ by $\tau(x_i) = \sigma(x_i)^{\mathbf{w}}$ for $i = 1, \dots, n$. Then, $\sigma(m)^{\mathbf{w}} = \tau(m)$ holds

for each monomial m by (2.1). Put $\alpha = \deg_{\mathbf{w}} f$ and write $f^{\mathbf{w}} = \sum_i m_i$, where m_i 's are nonzero monomials of $k[\mathbf{x}]_{\alpha}$. Then, we have $\deg_{\mathbf{w}} \sigma(m_i) = \deg_{\mathbf{w}} m_i = \alpha$ for each i , and $\sum_i \sigma(m_i)^{\mathbf{w}} = \sum_i \tau(m_i) = \tau(\sum_i m_i) = \tau(f^{\mathbf{w}}) \neq 0$. This implies that

$$\sum_i \sigma(m_i)^{\mathbf{w}} = \left(\sum_i \sigma(m_i) \right)^{\mathbf{w}} = \sigma \left(\sum_i m_i \right)^{\mathbf{w}} = \sigma(f^{\mathbf{w}})^{\mathbf{w}}.$$

Hence, $\sigma(f^{\mathbf{w}})^{\mathbf{w}}$ belongs to $k[\mathbf{x}]_{\alpha}$. Therefore, we get $\deg_{\mathbf{w}} \sigma(f^{\mathbf{w}}) = \deg_{\mathbf{w}} f$. $\square$

At this point, it is useful to remark that $\gamma < |\mathbf{w}|_*$ implies $\gamma \leq |\mathbf{w}| - w_{n-1}$ for each $\gamma \in \mathbf{N}w_1 + \cdots + \mathbf{N}w_{n-1}$, and $|\mathbf{w}|_* \leq |\mathbf{w}| + w_1 - w_{n-1}$ by the definition of $|\mathbf{w}|_*$. Under the assumption that $w_1 \leq \cdots \leq w_n$, we have the following lemma.

Lemma 4.7. *If $\deg_{\mathbf{w}} \omega < |\mathbf{w}|_*$, then $h_1, \dots, h_{n-1}$ belong to k , h_n belongs to $k[\mathbf{x} \setminus \{x_n\}]$ and $\deg_{\mathbf{w}} h_n \leq w_n - w_{n-1}$.*

Proof. We remark that $\deg_{\mathbf{w}} \eta_i = |\mathbf{w}| - w_i$ for $i = 1, \dots, n$. Hence, we have

$$\deg_{\mathbf{w}} h_i + |\mathbf{w}| - w_i = \deg_{\mathbf{w}} h_i \eta_i \leq \deg_{\mathbf{w}} \omega < |\mathbf{w}|_* \leq |\mathbf{w}| + w_1 - w_{n-1},$$

and so $\deg_{\mathbf{w}} h_i < w_1 - w_{n-1} + w_i$. Since $w_1 \leq \cdots \leq w_n$ by assumption, we know that $\deg_{\mathbf{w}} h_i < w_1$ for $1 \leq i \leq n-1$, and $\deg_{\mathbf{w}} h_n < w_n$. Thus, h_i belongs to k if $i < n$, and to $k[\mathbf{x} \setminus \{x_n\}]$ if $i = n$. Since $\deg_{\mathbf{w}} \eta_n = w_1 + \cdots + w_{n-1}$, it follows that $\deg_{\mathbf{w}} h_n \eta_n$ belongs to $\mathbf{N}w_1 + \cdots + \mathbf{N}w_{n-1}$. Since $\deg_{\mathbf{w}} h_n \eta_n \leq \deg_{\mathbf{w}} \omega < |\mathbf{w}|_*$, this implies that $\deg_{\mathbf{w}} h_n \eta_n \leq |\mathbf{w}| - w_{n-1}$ as remarked. Therefore, we have $\deg_{\mathbf{w}} h_n \leq w_n - w_{n-1}$. $\square$

Now, we prove Lemma 4.5. Let I be the set of $1 \leq i < n$ such that $h_i \neq 0$. When $I = \emptyset$, the assertion holds for $\phi = \text{id}_{k[\mathbf{x}]}$, $h = h_n$ and $l = n$. Hence, we may assume that $I \neq \emptyset$. We prove the assertion by induction on the number of elements of I . First, assume that $I = \{r\}$ for some $1 \leq r < n$. Then, we have $\omega = h_r \eta_r + h_n \eta_n$ with $h_r \neq 0$. Since $\deg_{\mathbf{w}} \omega < |\mathbf{w}|_*$ by assumption, we know by Lemma 4.7 that h_r belongs to $k^\times$, h_n belongs to $k[\mathbf{x} \setminus \{x_n\}]$ and $\deg_{\mathbf{w}} h_n \leq w_n - w_{n-1}$. By integrating $h_r^{-1} h_n$ with respect to x_r , we obtain $g \in k[\mathbf{x} \setminus \{x_n\}]$ such that $\partial g / \partial x_r = h_r^{-1} h_n$ and

$$\deg_{\mathbf{w}} g = \deg_{\mathbf{w}} h_n x_r = \deg_{\mathbf{w}} h_n + w_r \leq (w_n - w_{n-1}) + w_r \leq w_n.$$

Define $\phi \in \text{T}_n(k)$ by $\phi(x_n) = x_n + (-1)^{n-r} g$ and $\phi(x_i) = x_i$ for each $i \neq n$. Then, we have $\deg_{\mathbf{w}} \phi(f_i) = \deg_{\mathbf{w}} f_i$ for each i by Lemma 4.6 and the

preceding inequality. Since h_r and h_n are fixed under ϕ , we know by (4.1) and (4.2) that

$$\begin{aligned}\omega^\phi &= \phi(h_r)\eta_r^\phi + \phi(h_n)\eta_n^\phi = h_r\eta_r^\phi + h_n\eta_n = h_r\left(\eta_r - \frac{\partial g}{\partial x_r}\eta_n\right) + h_n\eta_n \\ &= h_r(\eta_r - h_r^{-1}h_n\eta_n) + h_n\eta_n = h_r\eta_r.\end{aligned}$$

Therefore, the assertion is true when $\#I = 1$.

Next, assume that $\#I \geq 2$. Let $r, s \in I$ be such that $1 \leq r < s < n$. Since $h_1, \dots, h_{n-1}$ belong to k by Lemma 4.7, we can define $\sigma \in T_n(k)$ by $\sigma(x_s) = x_s + (-1)^{s-r}h_r^{-1}h_sx_r$ and $\sigma(x_i) = x_i$ for each $i \neq s$. Then, $\deg_{\mathbf{w}}\sigma(f) = \deg_{\mathbf{w}}f$ holds for each $f \in k[\mathbf{x}]$ by Lemma 4.6, since $\deg_{\mathbf{w}}h_r^{-1}h_sx_r = w_r \leq w_s$. By (4.2), we have $\eta_r^\sigma = \eta_r - h_r^{-1}h_s\eta_s$ and $\eta_i^\sigma = \eta_i$ for each $i \neq r$. Put $f'_i = \sigma(f_i)$ for $i = 1, \dots, n-1$. Then, it follows that

$$\begin{aligned}df'_1 \wedge \dots \wedge df'_{n-1} &= \omega^\sigma = \sigma(h_r)\eta_r^\sigma + \sum_{i \neq r, n} \sigma(h_i)\eta_i^\sigma + \sigma(h_n)\eta_n^\sigma \\ &= h_r(\eta_r - h_r^{-1}h_s\eta_s) + \sum_{i \neq r, n} h_i\eta_i + \sigma(h_n)\eta_n = \sum_{i \neq s, n} h_i\eta_i + \sigma(h_n)\eta_n.\end{aligned}$$

Since $\deg_{\mathbf{w}}\sigma(h_n) = \deg_{\mathbf{w}}h_n$, this implies that

$$\deg_{\mathbf{w}}df'_1 \wedge \dots \wedge df'_{n-1} \leq \deg_{\mathbf{w}}\omega < |\mathbf{w}|_*.$$

Thus, by induction assumption, there exist $\tau \in T_n(k)$, $h \in k[\mathbf{x}]$ and $1 \leq l \leq n$ such that

$$d\tau(f'_1) \wedge \dots \wedge d\tau(f'_{n-1}) = h\eta_l$$

and $\deg_{\mathbf{w}}\tau(f'_i) = \deg_{\mathbf{w}}f'_i$ for $i = 1, \dots, n-1$. Then, $\phi := \tau \circ \sigma$ is a tame automorphism such that $\omega^\phi = h\eta_l$. Moreover, we have $\deg_{\mathbf{w}}\phi(f_i) = \deg_{\mathbf{w}}f_i$ for each i , since $\deg_{\mathbf{w}}\sigma(f_i) = \deg_{\mathbf{w}}f_i$. Therefore, the assertion is true when $\#I \geq 2$. This completes the proof of Lemma 4.5, and thereby completing the proof of Theorem 4.1.

5 Proof of Theorem 2.5

Throughout this section, we assume that $n = 3$. The goal of this section is to complete the proof of Theorem 2.5.

Lemma 5.1. *For any $\mathbf{w} \in (\Gamma_+)^3$ and $(d_1, d_2, d_3) \in \text{mdeg}_{\mathbf{w}}(\text{Aut}_k k[\mathbf{x}])$, the following assertions hold:*

- (i) (K1) and (K2) imply that at least two of d_1, d_2 and d_3 do not belong to $\{w_1, w_2, w_3\}$.
- (ii) (K1) and (K2) imply that $\Delta_{\mathbf{w}}(d_i, d_j) \geq |\mathbf{w}|_*$ for each $i, j \in \{1, 2, 3\}$ with $i \neq j$.
- (iii) If d_1 and d_2 are linearly dependent over $\mathbf{Z}$, then (K2) and (B1) imply (B2).
- (iv) If d_1 and d_2 are linearly dependent over $\mathbf{Z}$, then (K1), (K2), (A) and (B2) imply (K3), (K4) and (K5).
- (v) If d_1 and d_2 are linearly independent over $\mathbf{Z}$, then (K1), (K2) and (A) imply (K3) and (K4).

Proof. (i) Let $F \in \text{Aut}_k k[\mathbf{x}]$ be such that $\text{mdeg}_{\mathbf{w}} F = (d_1, d_2, d_3)$. By changing the indices of x_1, x_2 and x_3 , and w_1, w_2 and w_3 if necessary, we may assume that $w_1 \leq w_2 \leq w_3$. Then, we have $d_i \geq w_i$ for $i = 1, 2, 3$ by Lemma 2.1, since $d_1 < d_2 < d_3$ by (K1). First, we show that f_1 or f_2 does not belong to $k[x_1, x_2]$ by contradiction. Suppose that f_1 and f_2 both belong to $k[x_1, x_2]$. Then, $F' := (f_1, f_2)$ belongs to $\text{Aut}_k k[x_1, x_2]$, and $f_3 = \alpha x_3 + p$ for some $\alpha \in k^\times$ and $p \in k[x_1, x_2]$. If $(d_1, d_2) \neq (w_1, w_2)$, then we have $\deg_{\mathbf{w}} F' = d_1 + d_2 > w_1 + w_2$, since $d_i \geq w_i$ for each i . Hence, d_i belongs to $\mathbf{N}d_j$ for some $(i, j) \in \{(1, 2), (2, 1)\}$ by Proposition 4.2. Since $d_1 < d_2$, it follows that d_2 belongs to $\mathbf{N}d_1$, contradicting (K2). Assume that $(d_1, d_2) = (w_1, w_2)$. Then, we have $d_3 > w_3$, since $d_1 + d_2 + d_3 > |\mathbf{w}|$ by (K1). Hence, we get

$$d_3 = \deg_{\mathbf{w}} f_3 = \deg_{\mathbf{w}} (\alpha x_3 + p) = \deg_{\mathbf{w}} p.$$

Since p is an element of $k[x_1, x_2]$, it follows that d_3 belongs to $\mathbf{N}_0 w_1 + \mathbf{N}_0 w_2 = \mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$. This contradicts (K2). Therefore, f_1 or f_2 does not belong to $k[x_1, x_2]$. Since $w_1 \leq w_2 \leq w_3$ and $d_1 < d_2 < d_3$, this implies that $w_3 \leq d_2$. Now, suppose that (i) is false. Then, d_1 and d_2 must be at most w_3 . Hence, we have $d_2 = w_3$, and so $d_1 = w_l$ for some $l \in \{1, 2\}$. Thus, we may write $f_1 = \alpha_1 x_l + p_1$ and $f_2 = \alpha_2 x_3 + p_2$, where $\alpha_1, \alpha_2 \in k$, and $p_1 \in k[x_1, \dots, x_{l-1}]$ and $p_2 \in k[x_1, x_2]$ are such that $\deg_{\mathbf{w}} p_1 \leq w_l$ and $\deg_{\mathbf{w}} p_2 \leq w_3$. We note that $\alpha_2 \neq 0$, for otherwise f_1 and f_2 both belong to $k[x_1, x_2]$. If $\alpha_1 = 0$, then $f_1 = p_1$ belongs to $k[x_1, \dots, x_{l-1}]$. Since f_1 is not a constant, it follows that $l = 2$, and $f_1 = \alpha x_1 + \beta$ for some $\alpha \in k^\times$ and $\beta \in k$. Hence, we may assume that $\alpha_1 \neq 0$ by changing l if necessary. Since $k[x_l - p_1, x_m] = k[x_1, x_2]$ holds for $m \in \{1, 2\}$ with $m \neq l$, we can define $\sigma \in \text{Aut}_k k[x_1, x_2]$ by $\sigma(x_l) = \alpha_1^{-1}(x_l - p_1)$ and $\sigma(x_m) = x_m$. Then, σ preserves the $\mathbf{w}$ -degree of each element of $k[x_1, x_2]$ by Lemma 4.6. Since p_2 is an element of $k[x_1, x_2]$, we can extend σ to an element of $\text{Aut}_k k[\mathbf{x}]$ by setting $\sigma(x_3) = \alpha_2^{-1}(x_3 - \sigma(p_2))$. Then, σ also preserves the $\mathbf{w}$ -degree of each element of $k[\mathbf{x}]$, since $\deg_{\mathbf{w}} \sigma(p_2) = p_2 \leq w_3$. By definition, we have

$\sigma(f_1) = x_l$ and $\sigma(f_2) = x_3$, and so $\sigma(f_3) = \alpha x_m + p$ for some $\alpha \in k^\times$ and $p \in k[x_l, x_3]$. Since $\deg_{\mathbf{w}} x_m = w_m \leq w_3 = d_2 < d_3$, we get

$$d_3 = \deg_{\mathbf{w}} f_3 = \deg_{\mathbf{w}} \sigma(f_3) = \deg_{\mathbf{w}} (\alpha x_m + p) = \deg_{\mathbf{w}} p.$$

Therefore, d_3 belongs to $\mathbf{N}_0 w_l + \mathbf{N}_0 w_3 = \mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$, contradicting (K2).

(ii) Take any $i, j \in \{1, 2, 3\}$ with $i \neq j$. Then, d_i or d_j does not belong to $\{w_1, w_2, w_3\}$ by (i). Moreover, since $d_1 < d_2 < d_3$ by (K1), we see from (K2) that d_i and d_j do not belong to $\mathbf{N}d_j$ and $\mathbf{N}d_i$, respectively. Therefore, we have $\Delta_{\mathbf{w}}(d_i, d_j) \geq |\mathbf{w}|_*$ by the remark after Theorem 4.1.

(iii) Since d_1 and d_2 are linearly dependent over $\mathbf{Z}$ by assumption, we may write $d_i = u_i d$ for $i = 1, 2$, where $u_1, u_2 \in \mathbf{N}$ are such that $\gcd(u_1, u_2) = 1$, and $d = \gcd(d_1, d_2)$. Now, suppose that (B2) is not satisfied. Then, we have

$$d_3 \geq \text{lcm}(d_1, d_2) - d_1 - d_2 + |\mathbf{w}|_* = (u_1 u_2 - u_1 - u_2)d + |\mathbf{w}|_*.$$

Since $|\mathbf{w}|_* \geq d$ and $d_3 = u_3 d$ for some $u_3 \in \mathbf{N}$ by (B1), it follows that

$$u_3 \geq (u_1 u_2 - u_1 - u_2) + 1 = (u_1 - 1)(u_2 - 1).$$

Hence, u_3 belongs to $\mathbf{N}_0 u_1 + \mathbf{N}_0 u_2$ by Sylvester's formula for the Frobenius number. Therefore, d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$, contradicting (K2).

(iv) First, we check (K3) and (K4). By (ii), we have

$$\max\{\Delta_{\mathbf{w}}(d_i, d_j), |\mathbf{w}|_*\} = \Delta_{\mathbf{w}}(d_i, d_j) \quad (5.1)$$

for each $i \neq j$. Hence, (A) implies (K3) and (K4) if $3d_2 \neq 2d_3$ or $sd_1 \neq 2d_3$ for any odd number $s \geq 3$. Assume that $3d_2 = 2d_3$ and $sd_1 = 2d_3$ for some odd number $s \geq 3$. Then, there exists $d \in \Gamma_+$ such that $d_1 = 6d$, $d_2 = 2sd$ and $d_3 = 3sd$. Since d_2 does not belong to $\mathbf{N}d_1$ by (K2), we have $3 \nmid s$, and hence $\text{lcm}(d_1, d_2) = 6sd = 2d_3$. Thus, (B2) implies that

$$d_1 + d_2 < \text{lcm}(d_1, d_2) - d_3 + |\mathbf{w}|_* = d_3 + |\mathbf{w}|_* \leq d_3 + \Delta_{\mathbf{w}}(d_i, d_3)$$

for $i = 1, 2$ in view of (ii). Therefore, we get (K3) and (K4).

To show (K5), we may assume that $4d_1 = 3d_2$. Write $d_1 = 3d$ and $d_2 = 4d$, where $d = \gcd(d_1, d_2)$. Then, we have

$$d_3 < \text{lcm}(d_1, d_2) - d_1 - d_2 + |\mathbf{w}|_* = (3 \cdot 4 - 3 - 4)d + |\mathbf{w}|_* = 5d + |\mathbf{w}|_* \quad (5.2)$$

by (B2). Hence, it suffices to show that $|\mathbf{w}|_* \leq \Delta_{\mathbf{w}}(2d_1, d_2)$. By the remark after Theorem 4.1, we have only to check that $2d_1$ and d_2 do not belong to $\mathbf{N}d_2$ and $\mathbf{N}2d_1$, respectively, and that $2d_1$ or d_2 does not belong to $\{w_1, w_2, w_3\}$. Since $2d_1 = 6d$ and $d_2 = 4d$, the first part is clear. We prove the last part

by contradiction. Suppose that $2d_1$ and d_2 both belong to $\{w_1, w_2, w_3\}$. We may assume that $w_1 \leq w_2 \leq w_3$ as in the proof of (i). Then, $(2d_1, d_2)$ is equal to (w_2, w_1) or (w_3, w_2) , since $2d_1 = 6d > 4d = d_2$. We claim that the former case does not occur, for otherwise $w_i \geq w_1 = d_2 > d_1$ for any i , contradicting Lemma 2.1. Hence, we have $d_2 = w_2$. Since $d_1 < d_2$, this implies that f_1 belongs to $k[x_1]$, and hence is a linear polynomial in x_1 over k . Thus, we get $d_1 = w_1$. Therefore, d_1 and d_2 belong to $\{w_1, w_2, w_3\}$, contradicting (i).

(v) By (K1) and (K2), we have (5.1). Since d_1 and d_2 are linearly independent over $\mathbf{Z}$, we have $3d_2 \neq 2d_3$ or $sd_1 \neq 2d_3$ for any odd number $s \geq 3$. Hence, (A) implies (K3) and (K4). $\square$

Now, let us prove Theorem 2.5. Suppose to the contrary that there exists $F \in T_3(k)$ such that $\text{mdeg}_{\mathbf{w}} F = (d_1, d_2, d_3)$. First, assume that d_1 and d_2 are linearly independent over $\mathbf{Z}$. Then, we have (K3) and (K4) by Lemma 5.1 (v). Hence, (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$ by Theorem 2.2 (ii), a contradiction. Next, assume that d_1 and d_2 are linearly dependent over $\mathbf{Z}$. Then, we have (B1) or (B2). On account of Lemma 5.1 (iii), we may assume that (B2) is satisfied. Then, we get (K3), (K4) and (K5) by Lemma 5.1 (iv). Since F is tame by supposition, we have

$$\deg_{\mathbf{w}} df_1 \wedge df_2 \geq \Delta_{\mathbf{w}}(d_1, d_2) \geq |\mathbf{w}|_*$$

by Lemma 5.1 (ii). Hence, it follows from (B2) that

$$d_1 + d_2 + d_3 < \text{lcm}(d_1, d_2) + |\mathbf{w}|_* \leq \text{lcm}(d_1, d_2) + \deg_{\mathbf{w}} df_1 \wedge df_2.$$

Thus, F does not belong to $T_3(k)$ by Theorem 2.2 (i), a contradiction. Therefore, (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$. This completes the proof of Theorem 2.5.

6 Applications

In this section, we give applications of our theorems. Throughout, d_1, d_2 and d_3 denote positive integers. First, we briefly discuss when the assumptions of Theorem 1.3 are fulfilled. Set

$$d'_i = \frac{d_i}{\gcd(d_1, d_2, d_3)}$$

for $i = 1, 2, 3$. Clearly, d'_i is an odd number if so is d_i for each i .

In this notation, we have the following lemma.

Lemma 6.1. *Assume that $d_1, d_2, d_3 \in \mathbf{N}$ satisfy $d_1 \leq d_2 \leq d_3$ and (c). Then, we have (a) if one of the following conditions holds:*

- (1) d'_1 is an odd number, and d'_2 is an odd number or $d'_3 \not\equiv 0 \pmod{3}$.
- (2) $d_1 \neq 2 \gcd(d_1, d_3)$ and d'_2 is an odd number.
- (3) $d'_1 \equiv 0 \pmod{2^l}$ for some $l \geq 2$, and d'_2 and d'_3 are odd numbers.
- (4) d_3 is a prime number.
- (5) $d_3 - d_2 \geq d_1 - 2$.
- (6) d'_1 is an odd number, and $3d_2 \neq 2d_3$ or $2d_1 \leq d_2 + 5$.

Proof. It is easy to see that both (1) and (2) imply (a1). In the case of (3), we have $3d_2 \neq 2d_3$ since d'_2 is an odd number. Since $d'_1 \equiv 0 \pmod{2^l}$ for some $l \geq 2$, and d'_3 is an odd number, $sd_1 \neq 2d_3$ holds for any odd number s . Thus, we get (a1). We prove that (4) implies (a1) by contradiction. If $3d_2 = 2d_3$, then we have $d_2 = 2$. Since $d_1 \leq d_2 \leq d_3$, this implies that $d_1 = 1$ or $d_1 = d_2$, contradicting (c). If $sd_1 = 2d_3$ for some odd number $s \geq 3$, then we have $d_1 = 2$. Hence, d_2 and d_3 must be odd numbers in view of (c). Since $d_2 \leq d_3$, it follows that $d_3 = 2l + d_2 = ld_1 + d_2$ for some $l \in \mathbf{N}_0$, contradicting (c). Therefore, (4) implies (a1). Clearly, (5) is equivalent to (a2). Finally, we consider the case (6). Since d'_1 is an odd number, we have (a1) if $3d_2 \neq 2d_3$. So assume that $3d_2 = 2d_3$. Then, d_2 is an even number. Since $2d_1 \leq d_2 + 5$ by assumption, it follows that $2d_1 \leq d_2 + 4$. Thus, by noting $3d_2 = 2d_3$, we get $d_1 \leq d_2/2 + 2 = d_3 - d_2 + 2$. Therefore, (a2) is satisfied. $\square$

By means of Lemma 6.1, we get the following corollary to Theorem 1.3. Here, we note that the assumption (6.1) is fulfilled whenever $\gcd(d_1, d_2) = 1$.

Corollary 6.2. *Let $d_1, d_2, d_3 \in \mathbf{N}$ be such that $d_1 \leq d_2 \leq d_3$, and*

$$\gcd(d_1, d_2, d_3) = \gcd(d_1, d_2) \leq 3 \quad \text{or} \quad d_1 + d_2 + d_3 \leq \text{lcm}(d_1, d_2) + 2. \quad (6.1)$$

Assume that one of (1) through (6) in Lemma 6.1 is satisfied. Then, (d_1, d_2, d_3) belongs to $\text{mdeg } T_3(k)$ if and only if $d_1 \mid d_2$ or d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.

The “only if” part of Corollary 6.2 readily follows from Theorem 1.3, while the “if” part is due to Karaś [5, Proposition 2.2] (cf. Proposition 1.2).

In the following, we prove a variety of Karaś type theorems using our main results. The “if” parts of Corollaries 6.3 through 6.9 below follow from the proposition of Karaś as above. So we only check (the contrapositions of) the “only if” parts. Corollaries 6.3 through 6.7 are special cases of Corollary 6.2. We prove Corollaries 6.8, 6.9 and 6.10 using Theorem 1.3, and Corollary 6.11 using Theorem 2.5.

The following result is given by Karaś-Zygadło [10, Theorem 2.1]. This is a special case of (1) of Corollary 6.2.

Corollary 6.3 (Karaś-Zygadlo). *Let $d_3 \geq d_2 \geq d_1 \geq 3$ be integers such that d_1 and d_2 are odd numbers with $\gcd(d_1, d_2) = 1$. Then, (d_1, d_2, d_3) belongs to $\text{mdeg } T_3(k)$ if and only if d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.*

The following result is given by Karaś [7, Theorem 1.1].

Corollary 6.4 (Karaś). *Let $d_3 \geq d_2 \geq 3$ be integers. Then, $(3, d_2, d_3)$ belongs to $\text{mdeg } T_3(k)$ if and only if $3 \mid d_2$ or d_3 belongs to $\mathbf{N}_0 3 + \mathbf{N}_0 d_2$.*

This result of Karaś was generalized by Sun-Chen [20, Theorem 3.3] as follows. Here, we mention that Karaś [9, Theorem 7.1] independently obtained a similar result in which (iii) is replaced by the stronger assumption that $d_2 \geq 2d_1 - 3$.

Corollary 6.5 (Sun-Chen). *Let $d_3 \geq d_2 \geq d_1 \geq 3$ be integers with d_1 a prime number. Assume that one of the following conditions holds:*

- (i) $d_2 / \gcd(d_2, d_3) \neq 2$; (ii) $d_3 / \gcd(d_2, d_3) \neq 3$; (iii) $d_2 \geq 2d_1 - 5$.

Then, (d_1, d_2, d_3) belongs to $\text{mdeg } T_3(k)$ if and only if $d_1 \mid d_2$ or d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.

The “only if” part of Corollary 6.5 follows from (6) of Corollary 6.2. In fact, (i) or (ii) holds if and only if $3d_2 \neq 2d_3$, and d_1 is an odd prime number by assumption. Moreover, we have $\gcd(d_1, d_2) = 1$ if $d_1 \nmid d_2$.

The following result is given by Karaś [9, Theorems 6.2 and 6.10].

Corollary 6.6 (Karaś). (i) *Let d_2 and d_3 be odd numbers with $d_3 \geq d_2 \geq 5$. Then, $(4, d_2, d_3)$ belongs to $\text{mdeg } T_3(k)$ if and only if d_3 belongs to $\mathbf{N}_0 4 + \mathbf{N}_0 d_2$.*
(ii) *Let d_2 and d_3 be an odd number and an even number, respectively, such that $d_3 \geq d_2 \geq 5$ and $d_3 - d_2 \neq 1$. Then, $(4, d_2, d_3)$ belongs to $\text{mdeg } T_3(k)$ if and only if d_3 belongs to $\mathbf{N}_0 4 + \mathbf{N}_0 d_2$.*

In both (i) and (ii) of Corollary 6.6, we have $\gcd(4, d_2) = 1$ since d_2 is an odd number. Hence, (i) is a special case of (3) of Corollary 6.2. The “only if” part of (ii) follows from (5) of Corollary 6.2. In fact, if d_3 does not belong to $\mathbf{N}_0 4 + \mathbf{N}_0 d_2$, then we have $d_3 - d_2 \geq 2 = 4 - 2$, since $d_3 \geq d_2$ and $d_3 - d_2 \neq 1$ by assumption.

The following result is given by Li-Du [14, Theorems 3.3 and 4.2].

Corollary 6.7 (Li-Du). *Let $d_3 \geq d_2 \geq d_1 \geq 3$ be integers.*

(i) *When $d_1 / \gcd(d_1, d_3) \neq 2$ and d_2 is a prime number, (d_1, d_2, d_3) belongs to $\text{mdeg } T_3(k)$ if and only if $d_1 = d_2$ or d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.*

(ii) *When $\gcd(d_1, d_2) = 1$ and d_3 is a prime number, (d_1, d_2, d_3) belongs to $\text{mdeg } T_3(k)$ if and only if d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.*

The “only if” part of Corollary 6.7 (i) follows from (2) of Corollary 6.2, since d_2 is an odd prime number, and so $\gcd(d_1, d_2) = 1$ if $d_1 \neq d_2$. (ii) is a special case of (4) of Corollary 6.2.

The following result is given by Li-Du [15, Theorem 3.3].

Corollary 6.8 (Li-Du). *Let $a, d \in \mathbf{N}$ be such that $a \geq 3$. Assume that $4d \neq ta$ for any odd number $t \geq 1$. Then, $(a, a + d, a + 2d)$ belongs to $\text{mdeg } T_3(k)$ if and only if $a \mid 2d$.*

We can slightly extend this result as follows.

Corollary 6.9. *Let $a, d \in \mathbf{N}$ be such that $a \geq 3$. Assume that $a = 4l$ and $d = tl$ for some $l \in \mathbf{N}$ and an odd number $t \geq 1$ with $(t - 4)l + 2 \geq 0$. Then, $(a, a + d, a + 2d)$ belongs to $\text{mdeg } T_3(k)$ if and only if $a \mid 2d$.*

We prove the “only if” parts of Corollaries 6.8 and 6.9 together by means of Theorem 1.3. Set $(d_1, d_2, d_3) := (a, a + d, a + 2d)$. Assume that $a \nmid 2d$. Then, we have $a \nmid d$, and so $d_1 \nmid d_2$. We show that d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$ by contradiction. Suppose that $a + 2d = b_1 a + b_2(a + d)$ for some $b_1, b_2 \in \mathbf{N}_0$. Then, we have $b_2 \leq 1$, since a and d are positive. If $b_2 = 1$, then it follows that $d = b_1 a$, and hence $a \mid d$, a contradiction. If $b_2 = 0$, then we have $2d = (b_1 - 1)a$, and hence $a \mid 2d$, a contradiction. Thus, d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$. Therefore, (c) is satisfied. Next, we show (b2). Since $a \nmid d$, we have $a/\gcd(a, d) \geq 2$. We claim that $a/\gcd(a, d) \geq 3$. In fact, if $a = 2 \gcd(a, d)$, then a divides $2d$, a contradiction. Thus, we get

$$\text{lcm}(d_1, d_2) = \frac{d_1 d_2}{\gcd(d_1, d_2)} = \frac{a(a + d)}{\gcd(a, d)} \geq 3(a + d) = d_1 + d_2 + d_3.$$

This implies (b2). Finally, we show (a). In the case of Corollary 6.8, $sd_1 = sa$ is not equal to $2d_3 = 2(a + 2d)$ for any odd number $s \geq 3$, for otherwise $4d = (s - 2)a$. Since $a \neq d$, we get $3d_2 = 3(a + d) \neq 2(a + 2d) = 2d_3$. Thus, (a1) is satisfied. In the case of Corollary 6.9, we have

$$d_1 + d_2 = 4l + (4l + tl) \leq 8l + tl + \left((t - 4)l + 2\right) = 4l + 2tl + 2 = d_3 + 2.$$

Hence, (a1) is satisfied. Therefore, (d_1, d_2, d_3) does not belong to $\text{mdeg } T_3(k)$ by Theorem 1.3. This proves the “only if” parts of Corollaries 6.8 and 6.9.

The following corollary is a generalization of Karaš [9, Theorem 7.9] and Sun-Chen [20, Remark 3.9]. We mention that Karaš originally proved the case where d is a prime number with $5 \leq d \leq 31$, while Sun-Chen deduced from Corollary 6.5 that the assertion holds whenever d is a prime number with $d \geq 5$ (Sun-Chen also included the case where $d = 3$, but this is an obvious mistake in view of Proposition 1.2).

Corollary 6.10. *If $d \geq 5$ and $d \neq 6, 8$ for $d \in \mathbf{N}$, then $(d, 2(d-2), 3(d-2))$ does not belong to $\text{mdeg } T_3(k)$.*

Proof. Set $(d_1, d_2, d_3) := (d, 2(d-2), 3(d-2))$. We prove the assertion by means of Theorem 1.3. Since

$$d_1 + d_2 = 3d - 4 = 3(d-2) + 2 = d_3 + 2,$$

we have (a2). We check (c). Since $d \geq 5$, we see that $d_2 = 2(d-2)$ is not divisible by $d_1 = d$. We show that d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$ by contradiction. Suppose that $3(d-2) = a_1 d + a_2 \cdot 2(d-2)$ for some $a_1, a_2 \in \mathbf{N}_0$. Then, we have $a_2 \leq 1$, since $d \geq 5$. If $a_2 = 1$, then it follows that $(1 - a_1)d = 2$. This contradicts that a_1 and d are integers with $d \geq 5$. If $a_2 = 0$, then we have $(3 - a_1)d = 6$. Since $d \neq 6$ and $d \geq 5$, we get a contradiction similarly. Thus, d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$, proving (c). Finally, we show (b2). We claim that $d \geq 3 \gcd(d, 4)$. This is clear if $d \geq 12$. Since $d \geq 5$ and $d \neq 6, 8$, the rest of the cases are checked easily. Hence, we have

$$\text{lcm}(d_1, d_2) = \frac{d_1 d_2}{\gcd(d_1, d_2)} = \frac{d \cdot 2(d-2)}{\gcd(d, 4)} \geq 3 \cdot 2(d-2) = d_1 + d_2 + d_3 - 2,$$

proving (b2). Therefore, (d_1, d_2, d_3) does not belong to $\text{mdeg } T_3(k)$ thanks to Theorem 1.3. $\square$

The following result of Kanehira [4, Theorem 28] (cf. [1, Theorem 3.1]) is derived from Theorem 2.5.

Corollary 6.11 (Kanehira). *Let $d_3 \geq d_2 > d_1 \geq 3$ be integers such that d_1 and d_2 are odd numbers with $\gcd(d_1, d_2) = 1$. If there exist $\mathbf{w} \in \mathbf{N}^3$ and $F \in T_3(k)$ such that $\text{mdeg}_{\mathbf{w}} F = (d_1, d_2, d_3)$ and $\deg_{\mathbf{w}} F > |\mathbf{w}|$, then d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$.*

Proof. By assumption, (d_1, d_2, d_3) is an element of $\text{mdeg}_{\mathbf{w}} T_3(k)$ such that $d_1 < d_2 \leq d_3$ and $d_1 + d_2 + d_3 > |\mathbf{w}|$. Suppose to the contrary that d_3 does not belong to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$. Then, we have $d_2 \neq d_3$, and so $d_2 < d_3$. Thus, (K1) is satisfied. Since $d_1 \geq 3$ and $\gcd(d_1, d_2) = 1$, we see that d_2 does not belong to $\mathbf{N} d_1$. Hence, we get (K2) by supposition. Since d_1 and d_2 are odd numbers, (A1) is clear. Since $\gcd(d_1, d_2) = 1$, we see that (B1) is satisfied. Thus, (d_1, d_2, d_3) does not belong to $\text{mdeg}_{\mathbf{w}} T_3(k)$ by Theorem 2.5. This is a contradiction. Therefore, d_3 belongs to $\mathbf{N}_0 d_1 + \mathbf{N}_0 d_2$. $\square$

As noted after Theorem 4.1, $\Delta_{\mathbf{w}}(4, 6) \geq 4$ holds when $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, 1, 1)$ by Karaš [8, Theorem 14]. Modulo this result, Theorem 2.5 implies

Karaś [8, Theorem 1] asserting that $(4, 5, 6)$ does not belong to $\text{mdeg } T_3(k)$. In fact, $(d_1, d_2, d_3) = (4, 5, 6)$ satisfies (K1) and (K2) for $\Gamma = \mathbf{Z}$ and $\mathbf{w} = (1, 1, 1)$. Since $3d_1 = 12 = 2d_3$, and $d_1 + d_2 = 9$ is less than $d_3 + \Delta_{\mathbf{w}}(4, 6) = 10$, we see that (A3) holds for $s = 3$. Since $d_1 + d_2 + d_3 = 15$ is less than $\text{lcm}(d_1, d_2) = 20$, we get (B2).

References

- [1] E. Edo, T. Kanehira, M. Karaś and S. Kuroda, Separability of wild automorphisms of a polynomial ring, to appear in Transform. Groups.
- [2] A. van den Essen, Polynomial automorphisms and the Jacobian conjecture, Progress in Mathematics, Vol. 190, Birkhäuser, Basel, Boston, Berlin, 2000.
- [3] H. Jung, Über ganze birationale Transformationen der Ebene, J. Reine Angew. Math. **184** (1942), 161–174.
- [4] T. Kanehira, Weighted multidegrees of tame automorphisms of a polynomial ring in three variables (Japanese), Master’s Thesis, Tokyo Metropolitan University, February, 2012.
- [5] M. Karaś, There is no tame automorphism of $\mathbb{C}^3$ with multidegree $(3, 4, 5)$, Proc. Amer. Math. Soc. **139** (2011), 769–775.
- [6] M. Karaś, Tame automorphisms of $\mathbb{C}^3$ with multidegree of the form (p_1, p_2, d_3) , Bull. Pol. Acad. Sci. Math. **59** (2011), 27–32.
- [7] M. Karaś, Tame automorphisms of $\mathbb{C}^3$ with multidegree of the form $(3, d_2, d_3)$, J. Pure Appl. Algebra **214** (2010), 2144–2147.
- [8] M. Karaś, There is no tame automorphism of $\mathbb{C}^3$ with multidegree $(4, 5, 6)$, arXiv:math.AG/1104.1061v1.
- [9] M. Karaś, Multidegrees of tame automorphisms of $\mathbb{C}^n$, Dissertationes Math. (Rozprawy Mat.) **477** (2011), 55 pp.
- [10] M. Karaś and J. Zygałło, On multidegrees of tame and wild automorphisms of $\mathbf{C}^3$, to appear in J. Pure. Appl. Algebra.
- [11] S. Kuroda, A generalization of the Shestakov-Umirbaev inequality, J. Math. Soc. Japan **60** (2008), 495–510.

- [12] S. Kuroda, Shestakov-Umirbaev reductions and Nagata's conjecture on a polynomial automorphism, *Tohoku Math. J.* **62** (2010), 75–115.
- [13] S. Kuroda, Wildness of polynomial automorphisms in three variables, [arXiv:math.AC/1110.1466v1](https://arxiv.org/abs/math.AC/1110.1466v1).
- [14] J. Li and X. Du, Multidegrees of tame automorphisms with one prime number, [arXiv:math.AC/1204.0930](https://arxiv.org/abs/math.AC/1204.0930)
- [15] J. Li and X. Du, Tame automorphisms with multidegrees in the form of arithmetic progressions, [arXiv:math.AC/1112.6071v1](https://arxiv.org/abs/math.AC/1112.6071v1).
- [16] H. Matsumura, Commutative ring theory, translated from the Japanese by M. Reid, second edition, Cambridge Studies in Advanced Mathematics, 8, Cambridge Univ. Press, Cambridge, 1989.
- [17] M. Nagata, On Automorphism Group of $k[x, y]$, Lectures in Mathematics, Department of Mathematics, Kyoto University, Vol. 5, Kinokuniya Book-Store Co. Ltd., Tokyo, 1972.
- [18] I. Shestakov and U. Umirbaev, Poisson brackets and two-generated subalgebras of rings of polynomials, *J. Amer. Math. Soc.* **17** (2003), 181–196.
- [19] I. Shestakov and U. Umirbaev, The tame and the wild automorphisms of polynomial rings in three variables, *J. Amer. Math. Soc.* **17** (2003), 197–227.
- [20] X. Sun and Y. Chen, Multidegrees of Tame Automorphisms in Dimension Three, *Publ. Res. Inst. Math. Sci.* **48** (2012), 129–137.

Department of Mathematics and Information Sciences
Tokyo Metropolitan University
1-1 Minami-Osawa, Hachioji
Tokyo 192-0397, Japan
kuroda@tmu.ac.jp